



HAL
open science

A characterization of b-chromatic and partial Grundy numbers by induced subgraphs

Brice Effantin, Nicolas Gastineau, Olivier Togni

► **To cite this version:**

Brice Effantin, Nicolas Gastineau, Olivier Togni. A characterization of b-chromatic and partial Grundy numbers by induced subgraphs. 2015. hal-01157902v1

HAL Id: hal-01157902

<https://u-bourgogne.hal.science/hal-01157902v1>

Preprint submitted on 28 May 2015 (v1), last revised 28 Apr 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A characterization of b-chromatic and partial Grundy numbers by induced subgraphs

Brice Effantin², Nicolas Gastineau¹ and Olivier Togni¹

¹*Université de Bourgogne, 21078 Dijon cedex, France, Le2i, UMR CNRS 6303*

²*Université de Lyon, CNRS, Université Lyon 1, LIRIS, UMR5205, F-69622, France*

May 28, 2015

Abstract

Gyárfás et al. and Zaker have proven that the Grundy number of a graph G satisfies $\Gamma(G) \geq t$ if and only if G contains an induced subgraph called a t -atom. The family of t -atoms has bounded order and contains a finite number of graphs. In this article, we introduce equivalents of t -atoms for b-coloring and partial Grundy coloring. This concept is used to prove that determining if $\varphi(G) \geq t$ and $\partial\Gamma(G) \geq t$ (under conditions for the b-coloring), for a graph G , is in XP with parameter t . We illustrate the utility of the concept of t -atoms by giving results on b-critical vertices and edges, on b-perfect graphs and on graphs of girth at least 7.

1 Introduction

Given a graph G , a *proper k -coloring* of G is a surjective function $c : V(G) \rightarrow \{1, \dots, k\}$ such that $c(u) \neq c(v)$ for any $uv \in E(G)$; the *color class* V_i is the set $\{u \in V \mid c(u) = i\}$ and a vertex v has *color* i if $v \in V_i$. A vertex v of color i is a *Grundy vertex* if it is adjacent to at least one vertex colored j , for every $j < i$. A *Grundy k -coloring* is a proper k -coloring such that every vertex is a Grundy vertex. The *Grundy number* of a graph G , denoted by $\Gamma(G)$, is the largest integer k such that there exists a Grundy k -coloring of G [11]. A *partial Grundy k -coloring* is a proper k -coloring such that every color class contains at least one Grundy vertex. The *partial Grundy number* of a graph G , denoted by $\partial\Gamma(G)$, is the largest integer k such that there exists a partial Grundy k -coloring of G . Let G and G' be two graphs. By $G \cup G'$ we denote the graph with vertex set $V(G) \cup V(G')$ and edge set $E(G) \cup E(G')$. Let $m(G)$ be the largest integer m such that G has at least m vertices of degree at least $m - 1$. A graph G is *tight* if it has exactly $m(G)$ vertices of degree $m(G) - 1$.




Figure 1: The graph $K_{3,3}^-$ with $\varphi(K_{3,3}^-) = 2$ (on the left) and $\varphi_r(K_{3,3}^-) = 3$ (on the right).

Another coloring parameter with domination constraints on the colors is the *b-chromatic number*. In a proper- k -coloring, a vertex v of color i is a *b-vertex* if v is adjacent to at least one vertex colored j , $1 \leq j \neq i \leq k$. A *b- k -coloring*, also called *b-coloring* when k is not specified, is a proper k -coloring such that every color class contains at least one b-vertex. The *b-chromatic number* of a graph G , denoted by $\varphi(G)$, is the largest integer k such that there exists a b- k -coloring of G . In this paper, we introduce the concept of *b-relaxed number*, denoted by $\varphi_r(G)$. A *b- k -relaxed coloring* of G is a b- k -coloring of a subgraph of G . The b-relaxed number of G is $\varphi_r(G) = \max_{H \subseteq G} (\varphi(H))$, for H an induced subgraph of G . Note that we have $\varphi(G) \leq \varphi_r(G) \leq \partial\Gamma(G)$. The difference between $\varphi(G)$ and $\varphi_r(G)$ can be arbitrary large. Let $K_{n,n}^-$ denotes the complete bipartite graph $K_{n,n}$ in which we remove $n - 1$ pairwise non incident edges (or $n - 1$ edges of a perfect matching in $K_{n,n}$) [1]. For this graph we have $\varphi(K_{n,n}^-) = 2$ and $\varphi_r(K_{n,n}^-) = n$ as Figure 1 illustrates it (for $n = 3$).

The concept of b-coloring has been introduced by Irving and Manlove [17], and a large number of papers was published (see e.g. [8, 20]). The b-chromatic number of regular graphs has been investigated in a serie of papers ([6, 18, 21, 23]). Determining the b-chromatic number of a tight graph is NP-hard even for a connected bipartite graph [19] and a tight chordal graph [13].

In this paper, we study the decision problems b-COL, b-r-COL and pG-COL with parameter t from Table 1.

	b-COL	b-r-COL	G-COL	pG-COL
Question	Does $\varphi(G) \geq t$?	Does $\varphi_r(G) \geq t$?	Does $\Gamma(G) \geq t$?	Does $\partial\Gamma(G) \geq t$?
Complexity class	undetermined	XP	XP [24]	XP

Table 1: The different decision problems with input a graph G and parameter t and their complexity class.

A decision problem is in FPT with parameter t if there exists an algorithm which resolves the problem in time $O(f(t) n^c)$, for an instance of size n , a computable function f and a constant c . A decision problem is in XP with parameter t if

there exists an algorithm which resolves the problem in time $O(f(t) n^{g(t)})$, for an instance of size n and two computable functions f and g .

The concept of t -atom was introduced independently by Gyárfás et al. [12] and by Zaker [24]. The family of t -atoms is finite and the presence of a t -atom can be determined in polynomial time for a fixed t . The following definition is slightly different from the definitions of Gyárfás et al. or Zaker, insisting more on the construction of every t -atom (some t -atoms can not be obtained with the initial construction of Zaker).

Definition 1.1 ([24]). *The family of t -atoms is denoted by \mathcal{A}_t^Γ , for $t \geq 1$, and is defined by induction. The family \mathcal{A}_1^Γ only contains K_1 . A graph G is in \mathcal{A}_{t+1}^Γ if there exists a graph G' in \mathcal{A}_t^Γ and an integer m , $m \leq |V(G')|$, such that G is composed of G' and an independent set I_m of order m , adding edges between G' and I_m such that every vertex in G' is connected to at least one vertex in I_m .*

Moreover, in the following sections, we say that a graph G in a family of graphs \mathcal{F} is *minimal*, if no graphs of \mathcal{F} is a proper induced subgraph of G . For example, a minimal t -atom A is a t -atom for which there are no t -atoms which are induced in A other than itself.

Theorem 1 ([12, 24]). *For a given graph G , $\Gamma(G) \geq t$ if and only if G contains an induced minimal t -atom.*

In this paper we prove equivalent theorems for b -relaxed number and partial Grundy number. In contrast with the minimal t -atoms, we can not define the minimal t -atoms for b -coloring as the smallest graphs such that G satisfies $\varphi(G) = t$ (also called b -critical graphs).

The paper is organized as follows: Section 2 is devoted to the definition of t -atoms for the partial Grundy coloring. This concept allows us to prove that the partial Grundy coloring problem is in XP with parameter t . Section 3 is similar to Section 2 but for b -relaxed-coloring. Section 4 is devoted to the concept of b -critical vertices and edges. Section 5 is about b -perfect graphs. Finally, Section 6 deals with graphs for which the b -relaxed and the b -chromatic numbers are equal.

2 Partial-Grundy- t -atoms: t -atoms for partial Grundy coloring

We start this section with the definition of t -atoms for partial Grundy coloring.

Definition 2.1. *Let t be an integer with $t \geq 2$. The family $\mathcal{A}^{\partial\Gamma}(t)$ of partial-Grundy- t -atoms (pG - t -atoms, for short) is obtained as follows: we begin by taking an independent set of t vertices, denoted by $C(t)$, called the center of the pG - t -atom and associating a different integer of $\{1, \dots, t\}$ to each vertex of $C(t)$. Let c_k be the vertex of $C(t)$ with associated integer k , $1 \leq k \leq t$. Let k be an integer with $1 \leq k \leq t$. A ∂ - k -dominating operation, on a graph G containing $C(t)$, consists in adding a set D of ℓ vertices, with $0 \leq \ell \leq t - k$ and*




Figure 2: The minimal pG-2-atom (on the left) and the three minimal pG-3-atoms (the numbers are the colors of the vertices and the surrounded vertices form the centers).

associated integer k and adding edges between $G - \{c_k\}$ and $D \cup \{c_k\}$ such that every vertex of $\{c_{k+1}, \dots, c_t\}$ is adjacent to a vertex of $D \cup \{c_k\}$ and $D \cup \{c_k\}$ remains independent. Let $\mathcal{G}_1 = \{C(t)\}$. The family of graphs \mathcal{G}_i , for $2 \leq i \leq t$, is obtained from the graphs in \mathcal{G}_{i-1} by doing a ∂ - i -dominating operation on these graphs. Finally, set $\mathcal{A}^{\partial\Gamma}(t) = \mathcal{G}_t$.

Note that the associated integers induce a partial Grundy coloring of the pG- t -atom. These associated integers of the pG- t -atoms are not considered anymore when we use the notion of induced subgraph. Figure 2 illustrates several pG- t -atoms obtained using the previous construction.

Observation 2.2. For every graph G in $\mathcal{A}^{\partial\Gamma}(t)$, we have $|V(G)| \leq \frac{t(t-1)}{2}$.

Lemma 2.3. Let t and t' be two integers such that $1 \leq t' < t$. Every graph in $\mathcal{A}^{\partial\Gamma}(t)$ contains a graph in $\mathcal{A}^{\partial\Gamma}(t')$ as induced subgraph.

Proof. Every graph G in $\mathcal{A}^{\partial\Gamma}(t)$ contains a graph G' of $\mathcal{A}^{\partial\Gamma}(t')$: we can obtain G' by removing every vertex with an associated integer k in G , for $t' < k \leq t$, and by removing, afterwards, the vertices of G' not adjacent to the centers. \square

Note that the only minimal pG-2-atom is P_2 . The minimal pG-3-atoms are C_3 , P_4 and $P_2 \cup P_3$. These graphs are illustrated in Figure 2.

Theorem 2. Let G be a graph. For a graph G , we have $\partial\Gamma(G) \geq t$ if and only if G contains an induced minimal pG- t -atom.

Proof. Suppose that $\partial\Gamma(G) = t'$ with $t' \geq t$. Thus, there exists a partial Grundy coloring of G with t' colors. Let $u_1, \dots, u_{t'}$ be a set of Grundy vertices, each in a different color class of $V(G)$. The graph induced by $N[u_1] \cup \dots \cup N[u_{t'}]$ contains a pG- t' -atom. Hence, by Lemma 2.3, since G contains an induced pG- t' -atom, then it also contains an induced minimal pG- t -atom.

Suppose G contains an induced minimal pG- t -atom. Thus, the integers associated to the vertices correspond to a partial-Grundy coloring of this pG- t -atom. We can extend this coloring to a partial Grundy coloring of G with at least t colors, using a greedy algorithm [9] for the remaining vertices. \square

Proposition 2.4. *Let G be a graph of order n and let t be an integer. There exists an algorithm in time $O(n^{\frac{t(t-1)}{2}})$ to determine if $\partial\Gamma(G) \geq t$. Hence, the problem pG-COL with parameter t is in XP.*

Proof. By Theorem 2, it suffices to verify that G contains an induced minimal pG- t -atom to have $\partial\Gamma(G) \geq t$. Since the size of a minimal pG- t -atom is bounded by $\frac{t(t-1)}{2}$, we obtain an algorithm in time $O(n^{\frac{t(t-1)}{2}})$. \square

We finish this section by determining every graph G with $\partial\Gamma(G) = 2$.

Proposition 2.5. *For a graph G , we have $\partial\Gamma(G) = 2$ if and only if $G = K_{n,m}$, for some $n \geq 1$ and $m \geq 1$.*

Proof. Zaker [24] has proven that $\Gamma(G) = 2$ if and only if G is the disjoint union of copies of some $K_{n,m}$, for $n \geq 1$ and $m \geq 1$. Let n and m be positive integers. We can note that a graph containing two disjoint copies of $K_{n,m}$ contains an induced $P_2 \cup P_3$, hence a pG-3-atom. Hence, if $\partial\Gamma(G) = 2$, then $G = K_{n,m}$.

Moreover, $K_{n,m}$ does not contain induced C_3 , P_4 and $P_3 \cup P_2$. Hence, $\partial\Gamma(K_{n,m}) = 2$. \square

3 b- t -atoms: t -atoms for b-coloring

As in the previous section, we start this section with the definition of b- t -atoms (the notion of t -atom for b-coloring).

Definition 3.1. *Let t be an integer with $t \geq 2$. The family of b- t -atoms $\mathcal{A}^\varphi(t)$ is obtained as follows: we begin by taking an independent set of t vertices, denoted by $C(t)$, called the center of the b- t -atom and associating a different integer of $\{1, \dots, t\}$ to each vertex of $C(t)$. Let c_k be the vertex of $C(t)$ with associated integer k , $1 \leq k \leq t$. Let k be an integer with $1 \leq k \leq t$. A k -dominating operation on a graph G containing $C(t)$, consists in adding a set D of ℓ vertices, with $0 \leq \ell \leq t - 1$ and associated integer k and adding edges between $G - \{c_k\}$ and $D \cup \{c_k\}$ such that every vertex of $C(t) - \{c_k\}$ is adjacent to a vertex of $D \cup \{c_k\}$ and $D \cup \{c_k\}$ remains independent. Let $\mathcal{G}_1 = \{C(t)\}$. The family of graphs \mathcal{G}_i , for $2 \leq i \leq t$, is obtained from the graphs in \mathcal{G}_{i-1} by doing an i -dominating operation on these graphs. Finally, set $\mathcal{A}^\varphi(t) = \mathcal{G}_t$.*

For a fixed b- t -atom, the set of possible associated integers is called the set of b- t -atom colorings.

Note that the associated integers induce a b-coloring of the b- t -atom. These associated integers of the b- t -atom are not considered anymore when we use




Figure 3: The minimal b-2-atom (on the left) and the five minimal b-3-atoms.

the notion of induced subgraph. Figure 3 illustrates several b- t -atoms obtained using the previous construction.

Observation 3.2. For every graph G in $\mathcal{A}^\varphi(t)$, we have $|V(G)| \leq t^2$.

Lemma 3.3. Let t and t' be two integers such that $1 \leq t' < t$. Every graph in $\mathcal{A}^\varphi(t)$ contains a graph in $\mathcal{A}^\varphi(t')$ as induced subgraph.

Proof. Every graph G in $\mathcal{A}^\varphi(t)$ contains a graph G' in $\mathcal{A}^\varphi(t')$: we can obtain G' by removing every vertex with an associated integer k in G , for $t' < k \leq t$, and by removing, afterwards, the vertices not adjacent to the centers. \square

Note that the only minimal b-2-atom is P_2 . The minimal b-3-atoms are C_3 , P_5 , C_5 , $P_3 \cup P_4$ and $P_3 \cup P_3 \cup P_3$. These graphs are illustrated in Figure 3.

Observation 3.4. Every minimal pG- t -atom is an induced subgraph of a minimal b- t -atom or a minimal t -atom (an atom for the Grundy number).

Proposition 3.5. Let G be a graph. If $\varphi(G) \geq t$, then G contains an induced minimal b- t -atom.

Proof. Suppose that $\varphi(G) = t'$, with $t' \geq t$. Thus, there exists a b-coloring of G with t' colors. Let $u_1, \dots, u_{t'}$ be a set of b-vertices, each in a different color class of $V(G)$. The graph induced by $N[u_1] \cup \dots \cup N[u_{t'}]$ contains a b- t' -atom. Hence, by Lemma 3.3, since G contains an induced b- t' -atom, then it also contains an induced minimal b- t -atom. \square

Theorem 3. For a graph G , we have $\varphi_r(G) \geq t$ if and only if G contains an induced minimal b- t -atom.

Proof. Suppose that the graph G contains an induced b- t -atom A . Since A admits, by definition, a b- t -coloring, we have $\varphi_r(G) \geq t$. Using Proposition 3.5, we obtain the reciprocal. \square

Definition 3.6. Let G be a graph. For an induced subgraph A of G , let $N(A) = \{v \in V(G) \setminus V(A) \mid uv \in E(G), u \in V(A)\}$. A b - t -atom A is feasible in G if there exists a b - t -coloring of $V(A)$ that can be extended to the vertices of $N(A)$ without using new colors.

Proposition 3.7. Let G be a graph. If G contains an induced feasible minimal b - t -atom and no induced feasible minimal b - t' -atom, for $t' > t$, then $\varphi(G) = t$.

Proof. If G contains no induced minimal b - t -atom, then by Proposition 3.5, we have $\varphi(G) < t$. Suppose that G contains an induced feasible minimal b - t -atom A and no b - t -coloring of G exists. We begin by considering that the vertices of $A \cup N(A)$ are already colored with t colors. We can note that, by assumption, no coloring of $A \cup N(A)$ (from the definition) can be extended to the whole graph using only t colors. Let t' be the largest integer such that the coloring can not be extended to a b - t' -coloring of the whole graph and let v be a vertex that can not be given a color among $\{1, \dots, t'\}$. Thus, we suppose that the coloring can be extended to a b - $(t' + 1)$ -coloring where v is colored by $t' + 1$. Since $A \cup N(A)$ is already colored, we have $v \in V(G) \setminus (A \cup N(A))$. The vertex v should be adjacent to vertices of every color, otherwise it could be colored. One vertex of each color class in $N(v)$ should be adjacent to vertices of each color class (except its color). Otherwise, the colors of the vertices of $N(v)$ could be changed in order v be colored. Let $N_2[v]$ be the set of vertices at distance at most 2 from v . The set $N_2[v]$ induces a graph containing a b - t' -atom where $N[v]$ contains the center of this b - $(t' + 1)$ -atom. Moreover, this b - $(t' + 1)$ -atom is feasible as the whole graph is b - $(t' + 1)$ -colorable, contradicting the hypothesis. \square

Proposition 3.8. Let G be a graph. If $\varphi(G) = t$, then G contains an induced feasible minimal b - t -atom and no induced feasible minimal b - t' -atom, for $t' > t$.

Proof. Suppose $\varphi(G) = t$. By Proposition 3.5, G contains an induced minimal b - t -atom. If no induced minimal b - t -atom is feasible, then there exists no b - t -coloring of G , a contradiction. \square

A direct consequence of Proposition 3.7 and Proposition 3.8 is the following.

Theorem 4. For a graph G , we have $\varphi(G) = t$ if and only if G contains an induced feasible minimal b - t -atom and no induced feasible minimal b - t' -atom, for $t' > t$.

Note that the problem of determining if a graph has a b - t -coloring is NP-complete even if t is fixed [22]. However, it does not imply that determining if $\varphi(G) \geq t$ for a graph G is NP-complete. In contrast with the b -chromatic number, determining if a graph has b -relaxed number at least t is in XP.

Proposition 3.9. Let G be a graph of order n and let t be an integer. There exists an algorithm in time $O(n^{t^2})$ to determine if $\varphi_r(G) \geq t$. In particular, the problem b - r -COL with parameter t is in XP.

Proof. It suffices to verify that G contains an induced minimal b - t -atom to determine if $\varphi_r(G) \geq t$. By Observation 3.2, the size of a minimal b - t -atom is bounded by t^2 . Thus, we obtain an algorithm in time $O(n^{t^2})$. \square

Another NP-complete problem for a graph G is to determine its b -spectrum [2], i.e. the set of integers k such that G is b - k -colorable. For a graph G satisfying $\varphi(G) = \varphi_r(G)$, our algorithm can be used as the two concepts became the same.

Corollary 3.10. *Let G be a graph of order n and t be an integer. There exists an algorithm in time $O(n^{t^2})$ to determine if $\varphi(G) \geq t$ for a graph G satisfying $\varphi(G) = \varphi_r(G)$.*

4 b -critical vertices and edges

The concept of b -critical vertices and b -critical edges has been introduced recently and since five years a large number of articles are considering this subject [1, 4, 5, 10, 25]. In this section, we illustrate how this notion is strongly connected with the concept of b - t -atom.

Definition 4.1 ([4, 10]). *Let G be a graph. A vertex v of G is b -critical if $\varphi(G - v) < \varphi(G)$. An edge e is b -critical if $\varphi(G - e) < \varphi(G)$. A vertex v (edge e , respectively) in a graph G is a b - t -trap, if there exists a b - t -atom of G that becomes feasible by removing v (e , respectively).*

Proposition 4.2. *Let G be a graph. A vertex v is b -critical if and only if it is in every feasible minimal b - $\varphi(G)$ -atom and v is not a b - $\varphi(G)$ -trap.*

Proof. Let $t = \varphi(G)$. First, if v is not in a feasible minimal b - t -atom, then $\varphi(G - v) = t$ and v is not b -critical. If v is a b - t -trap, then, by definition, $\varphi(G - v) = t$. Second, suppose v is not a b - t -trap. If v is in every feasible minimal b - t -atom, then, since every minimal b - t -atom in G does not contain any other feasible minimal b - t -atom as induced subgraph, $G - v$ does not contain a feasible minimal b - t -atom. Thus, v is b -critical. \square

Corollary 4.3. *If a graph G contains two induced feasible minimal b - $\varphi(G)$ -atoms with disjoint set of vertices, then it contains no b -critical vertex.*

Corollary 4.4. *Let G be a graph and v be a vertex of $V(G)$. If $\varphi(G - v) > \varphi(G)$, then G contains a minimal b - t -atom which is not feasible, for some $t > \varphi(G)$. If $\varphi(G - v) < \varphi(G) - 1$, then $G - v$ contains no feasible minimal b - t -atom, for $\varphi(G - v) < t \leq \varphi(G)$.*

In [1], Balakrishnan and Raj have proved the following theorem.

Theorem 5 ([1]). *Let G be a graph and v be a vertex of $V(G)$. We have $\varphi(G) - \lfloor \frac{|V(G)|}{2} \rfloor + 2 \leq \varphi(G - v) \leq \varphi(G) + \lfloor \frac{|V(G)|}{2} \rfloor - 2$.*

Moreover, some families of graphs which satisfy properties from Theorem 5 are determined. More precisely, they have determined the families of graphs for which there exists a vertex v such that $\varphi(G - v) = \varphi(G) - \lfloor \frac{|V(G)|}{2} \rfloor + 2$ or $\varphi(G - v) = \varphi(G) + \lfloor \frac{|V(G)|}{2} \rfloor - 2$. In contrast with the b-chromatic number, we have the following property about the b-relaxed number.

Proposition 4.5. *Let G be a graph. If a vertex v is b-critical, then $\varphi_r(G - v) = \varphi_r(G) - 1$.*

Proof. By Proposition 4.2, v is in every $b\text{-}\varphi(G)$ -atom. Let i be the integer associated to v in the construction of this $b\text{-}\varphi(G)$ -atom. By removing the vertices with associated integer i , we obtain a $b\text{-}(\varphi(G) - 1)$ -atom and thus $\varphi_r(G - v) = \varphi_r(G) - 1$. \square

Note that this proposition was already proved for trees [4] (a tree T satisfies $\varphi(T) = \varphi_r(T)$, since every b-coloring of a subtree can be extended to the whole tree).

Lemma 4.6. *Let G be a graph with $4 \leq |V(G)| \leq 5$ and $E(G) \neq \emptyset$. We have $\varphi_r(G - v) = \varphi_r(G) + \lfloor \frac{|V(G)|}{2} \rfloor - 2$, for every vertex v of $V(G)$, if and only if G contains two disjoint edges but no induced minimal b-3-atom.*

Proof. We can note that we have $\varphi_r(G - v) = \varphi_r(G) + \lfloor \frac{|V(G)|}{2} \rfloor - 2$ if and only if $\varphi_r(G - v) = \varphi_r(G)$.

First, if G contains no minimal b-3-atom and contains an edge, then $\varphi_r(G) = 2$. Moreover, if G contains two disjoint edges, then for any vertex v , $G - v$ contains P_2 and $\varphi_r(G - v) = 2$.

Second, suppose that for every vertex v , $\varphi_r(G - v) = \varphi_r(G)$. The only minimal b-3-atoms that contains at most five vertices are K_3 , C_5 and P_5 . Moreover, the only minimal b-4-atoms and b-5-atoms that contain at most five vertices are K_4 and K_5 . We are going to show that G is not one of these graphs

Case 1: $\varphi_r(G) = 5$. If G is a K_5 , then, by removing any vertex v , we obtain $\varphi_r(G - v) = 4$.

Case 2: $\varphi_r(G) = 4$. If G is a K_4 , then, by removing any vertex v , we obtain $\varphi_r(G - v) = 3$. If G contains an induced K_4 , $|V(G)| = 5$ and G is not K_5 , then there exists a vertex v such $G - v$ has no induced K_4 and $\varphi_r(G - v) = 3$.

Case 3: $\varphi_r(G) = 3$. If G contains an induced K_3 and no induced K_4 , then, since the induced K_3 in G have a common vertex v , we obtain $\varphi_r(G - v) = 2$. Moreover, if G is P_5 or C_5 , then, by removing any vertex v , we obtain $\varphi_r(G - v) = 2$.

Thus, we can suppose that $\varphi_r(G) = 2$. If G contains only edges with a common vertex v , then $\varphi_r(G - v) = 1$. Hence, G contains no b-3-atom and contain two disjoint edges. \square

The following theorem is a generalization of a conjecture of Blidia et al. [3] for the parameter φ_r . Note that the graphs P_4 , C_4 and $P_2 \cup P_2$ do not contain any induced minimal b-3-atom and contain two disjoint edges.

Theorem 6. *Let G be a graph. We have $\varphi_r(G - v) = \varphi_r(G) + \lfloor \frac{|V(G)|}{2} \rfloor - 2$, for every vertex v of $V(G)$, if and only one of these conditions is true about G :*

- i) G is P_2 or C_3 .
- ii) $E(G) = \emptyset$ and $4 \leq |V(G)| \leq 5$.
- iii) $4 \leq |V(G)| \leq 5$ and G contains two disjoint edges but no b-3-atom.

Proof. Note that if $|V(G)| \geq 6$, then, by Proposition 4.5, we can not have $\varphi_r(G - v) < \varphi_r(G) + \lfloor |V(G)|/2 \rfloor - 2$. Note also that if G contains only one vertex, then it can not satisfy $\varphi_r(G - v) < \varphi_r(G) + \lfloor |V(G)|/2 \rfloor - 2$.

First, if $2 \leq |V(G)| \leq 3$, then we have $\varphi_r(G - v) = \varphi_r(G) - 1$ if and only if G is a minimal b- t -atom. Hence, if and only if G is P_2 or C_3 . Second, if G contains no edges, then $\varphi_r(G) = 1$ and for any vertex v , $\varphi_r(G - v) = 1$. The third condition is obtained by Lemma 4.6. \square

Definition 4.7. *Let t be a positive integer and A be a b- t -atom. An edge e is b-atom-critical in A if $A - e$ is not a b- t -atom.*

Proposition 4.8. *Let G be a graph. An edge e is b-critical if and only if it is b-atom-critical in every feasible minimal b- $\varphi(G)$ -atom and e is not a b- $\varphi(G)$ -trap.*

Proof. Let $t = \varphi(G)$. First, if e is not b-atom-critical in a feasible minimal b- t -atom, then $G - e$ contains a feasible minimal b- t -atom and $\varphi(G - e) = t$. If e is a b- t -trap, then, by definition, $\varphi(G - e) = t$. Second, suppose that e is not a b- t -trap. If e is b-atom-critical in every feasible minimal b- t -atom, then, since every feasible minimal b- t -atom in G does not contain any other feasible minimal b- t -atom as subgraph in $G - e$, the graph $G - e$ does not contain a feasible minimal b- t -atom. Thus, e is b-critical. \square

Corollary 4.9. *If a graph G contains two induced feasible minimal b- $\varphi(G)$ -atoms with disjoint sets of b-atom-critical edges, then G contains no b-critical edge.*

5 b-perfect graphs

A b-perfect graph is a graph for which every induced subgraph satisfies that its b-chromatic number is equal to its chromatic number. More generally, we present the following definitions.

Definition 5.1 ([14]). *A graph G is b- χ - k -bounded, for k a positive integer, if $\varphi(G') - \chi(G') \leq k$, for every induced subgraph G' of G . A b- χ -0-bounded graph is called a b-perfect graph. A χ - k -unbounded b-atom, for k a positive integer,*




Figure 4: The family \mathcal{F} : the imperfect b-atoms [15].

is a b-atom which is not b - χ - k -bounded and which does not contain any b - χ - k -bounded induced minimal b-atom other than itself. An imperfect b-atom is a b-atom which is not b-perfect and which does not contain any induced minimal b-atom which is not b-perfect other than itself.

Hoang et al. [15] characterized b-perfect graphs by giving the family \mathcal{F} of forbidden induced subgraphs depicted in Figure 4. We recall the following theorem:

Theorem 7 ([15]). *A graph is b-perfect if and only if it contains no graph from \mathcal{F} as induced subgraph.*

Note that every graph in the family \mathcal{F} is a b- t -atom for some t . More precisely, F_1 , F_2 and F_3 are the only minimal bipartite b-3-atoms. The remaining graphs are minimal b-4-atoms that do not contain F_1 , F_2 and F_3 as induced subgraph and which admit a proper coloring with three colors (as mentioned in [16]). We can state the following property about b- t -atoms.

Observation 5.2. *Let t and t' be two integers with $t' < t$. If a minimal b- t -atom A contains an induced χ - k -unbounded b- t' -atom, then A is not b - χ - k -bounded.*

Theorem 8. *Let k be a positive integer. A graph G is not b - χ - k -bounded if and only if it contains a χ - k -unbounded b -atom.*

Proof. First, if G contains a χ - k -unbounded b -atom, then, by definition, G is not χ - k -bounded.

Second, suppose G is not b - χ - k -bounded. Then, there exists an induced subgraph A of G of minimal order which is not b - χ - k -bounded. By removing vertices of A we can only decrease the chromatic number. Thus, by removing vertices we can obtain a b - $\varphi(A)$ -atom which is χ - k -unbounded. \square

Corollary 5.3. *The graphs with b -chromatic number t which are b - χ - k -bounded, for fixed integers k and t , can be defined by forbidding a finite family of induced subgraphs: the χ - k -unbounded b -atoms. Hence, a graph G is b -perfect if and only if it does not contain imperfect b -atoms.*

Let b - χ -BOUNDED be the following decision problem with parameter $\varphi(G)$ and let k be an integer, with $0 \leq k < \varphi(G)$.

b - χ - k -BOUNDED

Instance : A graph G .

Parameter : $\varphi(G)$.

Question: Does $\varphi(G) - \chi(G) \geq k$?

By Corollary 5.3, we obtain the following corollary:

Corollary 5.4. *Let G be a graph and k be an integer, with $0 \leq k < \varphi(G)$. There exists an algorithm in time $O(n^{\varphi(G)^2})$ to solve b - χ - k -BOUNDED.*

Since a graph G is b -perfect if and only if it does not contain imperfect b -atoms, we have the following theorem:

Theorem 9. *The number of imperfect b -atoms is finite. A graph is an imperfect b -atom if and only if it is in the family \mathcal{F} (Figure 4).*

The previous theorem is a consequence of Theorem 7. By proving that every b -4-atom except K_4 contains an induced subgraph of the family \mathcal{F} and by using Proposition 5.2, we can possibly obtain another proof of Theorem 7.

6 b -chromatic and b -relaxed chromatic numbers

In this section we consider the b -relaxed number relatively to the b -chromatic number. About the relation between these parameters we can state the following properties.

Proposition 6.1. *For a graph G satisfying $\varphi(G) = \Delta(G) + 1$, we have $\varphi(G) = \varphi_r(G)$.*

Proof. Since every vertex is adjacent to at most $\Delta(G)$ neighbors, it is always possible to extend a $(\Delta + 1)$ -coloring to a new vertex. \square

Proposition 6.2. For a tree T , we have $\varphi(T) = \varphi_r(T)$.

Proof. It is always possible to extend a k -coloring to a new vertex in a tree. \square

We finish this paper by proving that when a graph G has sufficiently large girth, we have $\varphi(G) = \varphi_r(G)$.

Theorem 10. Let G be a graph with girth g and $\varphi_r(G) \geq 3$. If $g \geq 7$, then $\varphi(G) = \varphi_r(G)$.

Proof. Suppose that $\varphi_r(G) = t$. Let A_t be a colored b - t -atom of G and let c_1, \dots, c_t be the vertices of the center of this b - t -atom. Moreover, we suppose that c_i is colored by i . If A_t is not feasible, then there exists a subset of vertices in $N(A_t)$ such that this coloring can not be extended to these vertices. By assumption, we suppose that A_t is not feasible. For a vertex $u \in N(A_t)$, we denote by $N_c(u)$ the set $\{i \in \{1, \dots, t\} \mid \exists v \in N(u) \cap N[c_i]\}$. We denote by $N[A_t]$, the set of vertices $A_t \cup N(A_t)$. In order to have a contradiction we begin by proving the following properties:

- i) if $u \in N(A_t)$, then u is not adjacent to two vertices of $N[c_i]$, for $1 \leq i \leq t$;
 - ii) if $u, v \in N(A_t)$ and $N_c(u) \cap N_c(v) \neq \emptyset$, then u and v are at distance at least 3;
 - iii) if $u, v \in N[c_i]$ and $u', v' \in N[c_i]$ for some i , $1 \leq i \leq t$, then no two pairs of vertices among $\{u, v, u', v'\}$ are adjacent.
- i) If u is adjacent to two vertices of $N[c_i]$, for some i , $1 \leq i \leq t$, then u is in a cycle of length at most 4.
 - ii) If two vertices u and v from $N(A_t)$ are at distance at most 2 and are both adjacent to vertices of $N[c_i]$, for $1 \leq i \leq t$, then u and v are in a cycle of length at most 6.
 - iii) If two pairs of vertices among $\{u, v, u', v'\}$ are adjacent then there is a cycle of length at most 6 in G .

We are going to prove that either each vertex $u \in N(A_t)$ can be given a color from $\{1, \dots, t\}$ or the graph contains a b - $(t+1)$ -atom (which contradicts $\varphi_r(G) = t$). First, we will consider the vertices from $N(A_t)$ which are not only adjacent to centers. Afterward, we will consider the remaining vertices of $N(A_t)$.

Case 1: u has at least one neighbor in A_t which is not a center.

By properties i) and ii), u has at most t neighbors in $N[A_t]$. If u has less than t neighbors in $N[A_t]$, then it can be given a color, hence assume u has exactly t neighbors in $N[A_t]$. If u has less than t neighbors in A_t , then there exists a neighbor w of u in $N(A_t)$. By property ii), $|N_c(w)| = 1$, i.e. w has two neighbors in $N[A_t]$ and thus can be recolored in order to give its original color to u . If u has exactly t neighbors in A_t , let v_1, \dots, v_ℓ be

the neighbors of u in A_t which are not in the center. Let c' be the color of v_1 and let c'' be the color of v_2 if it exists. Moreover, let i_1 and i_2 be two integers such that $v_1 \in N(c_{i_1})$ and $v_2 \in N(c_{i_2})$ (if $\ell \geq 2$).

Subcase 1.a: suppose that $\ell = 1$, i.e., u is adjacent to $t - 1$ vertices from the center and to one vertex in A_t which is not a center.

We can note that no vertex from $N[c_{i_1}]$ can have a neighbor in A_t , since it would create a cycle of length at most 6. Consequently, we can exchange the color of v_1 with the color of one vertex from $N(c_{i_1})$ and color u by c' .

Subcase 1.b: suppose that $\ell \geq 2$, i.e., u is adjacent to at most $t - 2$ vertices from the center and to at least two vertices in A_t which are not centers.

The vertex v_1 has no neighbor in $A_t - N[c_{i_1}]$ and v_2 has no neighbor in $A_t - N[c_{i_2}]$ since it would create a cycle of length at most 6. If a vertex of $N(c_{i_1}) - \{v_1\}$ has no neighbor of color c' , then we exchange the color of v_1 with the color of this vertex and color u by c' . If a vertex of $N(c_{i_2}) - \{v_2\}$ does not have a neighbor of color c'' , then we exchange the color of v_2 with the color of this vertex and color u by c'' . Thus, we suppose that every vertex of $N(c_{i_1}) - \{v_1\}$ has a neighbor of color c' and every vertex of $N(c_{i_2}) - \{v_2\}$ has a neighbor of color c'' .

In the case $c'' = i_1$ and $c' = i_2$, by property iii), no vertex of $N[c_{i_1}] \cup N[c_{i_2}]$ has a neighbor in $A_t - N[c_{i_1}] \cup N[c_{i_2}]$ other than these vertices of color c' and c'' . However, there could exist two adjacent vertices w and w' with $w \in N[c_{i_1}]$ and $w' \in N[c_{i_2}]$. If $t = 3$, then w and w' are colored by the same color and thus can not be adjacent. Consequently, we can exchange the color of v_1 with the color of v_2 , the color of c_{i_1} with the color of c_{i_2} and afterward we can exchange the color of one vertex from $N(c_{i_1}) - \{w, v_1\}$ (if $t = 3$, then we take w) with the color of v_1 and color u by c'' .

In the case $c'' \neq i_1$ (or $c' \neq i_2$, respectively), we suppose, without loss of generality, that $c' = i_2$ (or $c'' = i_1$, respectively). In this case, we exchange the color of c_{i_1} with the color of c_{i_2} and the color of v_1 with the color of a vertex w of color i_1 in $N(c_{i_2})$ (the color of v_2 with the color of a vertex w of color i_2 in $N(c_{i_1})$, respectively) and color u by c' (by c'' , respectively). We can note that the vertex w in $N(c_{i_2})$ (the vertex w in $N(c_{i_1})$, respectively) can be adjacent to a vertex of color c' (of color c'' , respectively). However, if w is adjacent to a vertex of color c' (of color c'' , respectively), then no vertex of $N(c_{i_2}) - \{w, v_2\}$ (of $N(c_{i_1}) - \{w, v_1\}$, respectively) is adjacent to a vertex of color c' (of color c'' , respectively). If w is adjacent to a vertex of this color, then we exchange the color of w with the color of a vertex of $N(c_{i_2}) - \{w, v_2\}$ (of $N(c_{i_1}) - \{w, v_1\}$, respectively). We can remark that if $t = 3$, then w can not be adjacent to a vertex of color c' (of color c'' , respectively) and the previous case never happen.

Case 2: every neighbor of u is in the center.

If u does not have neighbors in each color class, then we can give a color to u . The vertices from $N(u) \cap A_t$ are centers, and thus these vertices

have neighbors from each color class. If every vertex v in $N(u) \cap N(A_t)$ colored by a given color c does not have neighbors from each color class, then we recolor these vertices and give the color c to u . If u remains uncolorable, the set $N_2[u]$ induces a graph containing a $b-(t+1)$ -atom, which contradicts $\varphi_r(G) = t$.

□

We think that the previous theorem can be used to determine the family of graphs of girth at least 7 satisfying $\varphi(G) = m(G)$. It has already been proven that graphs of girth at least 7 have b -chromatic number at least $m(G) - 1$ [7].

Corollary 6.3. *Let G be a graph of girth at least 7 and of order n and let t be an integer. There exists an algorithm in time $O(n^{t^2})$ to determine if $\varphi(G) \geq t$.*

7 Open questions

We conclude this article by listing few open questions.

1. For which family of graphs are the b -relaxed number and the b -chromatic number equal?
2. Does there exist an easy characterization of feasible b - t -atoms?
3. Does there exist an FPT algorithm, with parameter t , to determine if $\varphi(G) \geq t$?

References

- [1] R. Balakrishnan and S. F. Raj. Bounds for the b -chromatic number of $G - v$. *Discrete Applied Mathematics*, 161:1173–1179, 2013.
- [2] D. Barth, J. Cohen, and T. Faik. On the b -continuity property of graphs. *Discrete Applied Mathematics*, 155:1761–1768, 2007.
- [3] M. Blidia, N. I. Eschouf, and F. Maffray. b -coloring of some bipartite graphs. *Australasian Journal of Combinatorics*, 53:67–76, 2012.
- [4] M. Blidia, N. I. Eschouf, and F. Maffray. On vertex b -critical tree. *Opuscula Math.*, 33:19–28, 2013.
- [5] M. Blidia, N. I. Eschouf, and F. Maffray. On edge b -critical graphs. *Discrete Applied Mathematics*, 180:176–180, 2015.
- [6] S. Cabello and M. Jakovac. On the b -chromatic number of regular graphs. *Discrete Applied Mathematics*, 159:1303–1310, 2011.
- [7] V. Campos, C. Lima, and A. Silva. Graphs of girth at least 7 have high b -chromatic number. *European Journal of Combinatorics*, 48:154–164, 2015.

- [8] B. Effantin and H. Kheddouci. The b-chromatic number of some power graphs. *Discrete Mathematics and Theoretical Computer Science*, 6:45–54, 2003.
- [9] P. Erdős, S. T. Hedetniemi, R. C. Laskar, and G. C. Prins. On the equality of the partial Grundy and upper chromatic number of graphs. *Discrete Mathematics*, 272:53–64, 2003.
- [10] N. I. Eschouf. Characterization of some b-chromatic edge critical graphs. *Australasian Journal of Combinatorics*, 47:21–35, 2010.
- [11] P. M. Grundy. Mathematics and games. *Eureka*, 2:6–8, 1939.
- [12] A. Gyárfás, Z. Király, and J. Lehel. On-line 3-chromatic graphs – ii critical graphs. *Discrete Mathematics*, 177:99–122, 1997.
- [13] F. Havet, C. L. Sales, and L. Sampaio. b-coloring of tight graphs. *Discrete Applied Mathematics*, 160:2709–2715, 2012.
- [14] C. T. Hoàng and M. Kouider. On the b-dominating coloring of graphs. *Discrete Applied Mathematics*, 152:4399–4408, 2005.
- [15] C. T. Hoàng, F. Maffray, and M. Mechebbek. A characterization of b-perfect graphs. *Journal of Graph Theory*, 71:95–122, 2012.
- [16] C. T. Hoàng, C. L. Sales, and F. Maffray. On minimally b-imperfect graphs. *Discrete Applied Mathematics*, 157:3519–3530, 2009.
- [17] R. W. Irving and D. F. Manlove. The b-chromatic number of a graph. *Discrete Applied Mathematics*, 91:127–141, 1999.
- [18] M. Jakovac and S. Klavžar. The b-Chromatic Number of Cubic Graphs. *Graphs and Combinatorics*, 26:107–118, 2010.
- [19] J. Kratochvíl, Z. Tuza, and M. Voigt. On the b-chromatic number of graphs. *Revised Papers from the 28th International Workshop on Graph-Theoretic Concepts in Computer Science*, pages 310–320, 2002.
- [20] F. Maffray and A. Silva. b-colouring the cartesian product of trees and some other graphs. *Discrete Applied Mathematics*, 161:650–669, 2013.
- [21] A. E. Sahili and M. Kouidder. b-chromatic number of regular graphs. *Utilitas Mathematica*, 80:211–216, 2009.
- [22] L. Sampaio. *Algorithmic aspects of graph colouring heuristics*. PhD thesis, University of Nice-Sophia Antipolis, 2012.
- [23] S. Shaebani. On the b-chromatic number of regular graphs without 4-cycle. *Discrete Applied Mathematics*, 160:1610–1614, 2012.
- [24] M. Zaker. Results on the Grundy chromatic number of graphs. *Discrete Mathematics*, 306:3166–3173, 2006.

- [25] M. Zamime, M. Kouider, and H. A. Haddadène. On the b-coloring of $G - e$. *European Journal of Combinatorics*, Available online, 2015.