



Inactivation of dried spores of *Bacillus subtilis* 168 by a treatment combining high temperature and pressure

Julia Hauck-Tiburski, Amauri Rosenthal, Cyril Iaconelli, Jean-Marie Perrier-Cornet, Patrick Gervais

► To cite this version:

Julia Hauck-Tiburski, Amauri Rosenthal, Cyril Iaconelli, Jean-Marie Perrier-Cornet, Patrick Gervais. Inactivation of dried spores of *Bacillus subtilis* 168 by a treatment combining high temperature and pressure. *International Journal of Food Microbiology*, 2019, 295, pp.1-7. 10.1016/j.ijfoodmicro.2019.01.017 . hal-02014283

HAL Id: hal-02014283

<https://u-bourgogne.hal.science/hal-02014283>

Submitted on 22 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 Inactivation of dried spores of *Bacillus subtilis* 168 by a treatment combining high
2 temperature and pressure

3 Julia Hauck-Tiburski^{a1}, Amauri Rosenthal^{a,b}, Cyril Iaconelli^a, Jean-Marie Perrier-Cornet^{a*}, Patrick
4 Gervais^a

5

6 a UMR Procédés Alimentaires et Microbiologiques (équipe PMB), AgroSup Dijon,
7 Université de Bourgogne Franche-Comté, 1, Esplanade Erasme, 21000 Dijon, France.

8 b Embrapa Food Technology, Av. das Américas, 29501 Guaratiba CEP 23020-470 Rio
9 de Janeiro - RJ, Brazil.

10 *: Corresponding author

11 e-mail:

12 Julia Hauck-Tiburski juliaha.dijon@gmail.com

13 Amauri Rosenthal amauri.rosenthal@embrapa.br

14 Cyril Iaconelli cyril.iaconelli@yahoo.fr

15 Jean-Marie Perrier-Cornet jperrier@u-bourgogne.fr

16 Patrick Gervais gervais@u-bourgogne.fr

17

¹ 'Present address': UFRRJ-DTA, Rodovia BR 465, km 7, CEP 23890-000 Seropédica, RJ, Brazil

18 Abstract

19 Specific treatments combining high temperatures of up to 150°C and moderate pressure of up
20 to 0.6 MPa have been applied to *Bacillus subtilis* 168 spores conditioned at different a_w levels
21 (between 0.10 and 0.70) corresponding to different residual water contents within the spore
22 core. The spores were treated as a dry powder in a pressurized nitrogen environment or in
23 water/glycerol solutions.

24 These thermodynamic conditions were intended to prevent any water evaporation from the
25 spore core during time/temperature treatments.

26 Our results clearly show that retaining liquid water in the core by applying pressure during the
27 treatment resulted in greater spore destruction (between 2.4 and 4.9 log at 150°C, 120 s and a_w
28 0.5 in powder) than the destruction observed after the treatment at atmospheric pressure (0.7
29 log), during which the water rapidly evaporated because its boiling point was reached.

30 Moreover, we found that the water activity level of the spore had a significant impact on spore
31 destruction: the higher the a_w level, the greater the spore inactivation.

32 We obtained similar results from spores heat-treated in powder and in water/glycerol solution
33 at the same a_w , confirming the strong influence of this parameter. We hypothesized that the
34 increased spore inactivation was related to the well-known thermal sensitivity of vital organic
35 molecules such as proteins, enzymes, and ribosomes in the presence of water.

36

37 Keywords

38 *Bacillus* spores – dry food – food powder - thermal sterilization – pressure – water activity

39

1. Introduction

Low water activity foods ($a_w < 0.6$), such as food powders, seeds, dried fruits, cakes, flakes, spices and aromatic herbs, represent a large amount of food products consumed directly or used as intermediate food products (Cuq et al., 2011). These products contain vegetative and sporulated bacteria that cannot grow due to the low water activity level. Safety problems can arise after rehydration before or during ingestion. To prevent such incidents, these products must be decontaminated, but heat decontamination is more difficult for dry products than it is for liquid products. Indeed, dried microorganisms are more heat-resistant (Grasso et al., 2014; Laroche and Gervais, 2003) and heating without water could also rapidly cause burning damage to the product. Among dried microorganisms, sporulated bacteria are particularly resistant (Tiburski et al., 2014) and spores of *Bacillus* species can survive various drying and heat treatments used in the food industry (Andersson et al., 1995). These spores have also been seen to adhere to stainless steel and to resist cleaning in place (CIP) procedures in and around food factories (Tauveron et al., 2006). These characteristics have obvious implications, from equipment bacteria could contaminate food and therefore cause high numbers of spores to be found in food products. Dried foods such as spices, milk powders and cereal products are often quite heavily contaminated with spores, and when water becomes available during food preparation these spores may germinate, leading to spoilage or food poisoning (Logan, 2012). However, destroying bacterial spores is rather difficult, and most of the techniques commonly used to treat dry foods result in very low spore inactivation (Silva et al., 2013).

The most frequently used decontamination techniques for dry products are steam thermal processing, irradiation and fumigation. However, steam adds undesirable moisture to dry foods and causes color changes. Fumigation has been banned in most countries due to its toxicity, and irradiation is strongly rejected by consumers. In addition, these decontamination

65 techniques cause loss of quality and sometimes low inactivation rates (Ghisleni et al., 2016).
66 Therefore, there is a need to develop new technologies to produce high quality and safe dry
67 food ingredients such as herbs and spices.

68 Though spices are non-perishable ingredients, once they are put in water-rich food products,
69 their natural or contaminant flora can quickly develop and multiply, consequently putting the
70 health of consumers at risk. This is an especially important issue for spices added to ready-to-
71 eat foods that are not subjected to further heat treatments (Van Doren et al., 2013).

72 Moreover the existence of pathogenic spores like *B. cereus* strains, that may cause foodborne
73 infections (Glasset et al., 2016) or like *Clostridium botulinum* strains that may sometimes
74 cause infant infection through dairy products (Doyle et al., 2015), is considered as a major
75 risk for the food industry.

76 It is thus important to decontaminate food powders efficiently, especially if these powders are
77 to be incorporated into a more complex preparation with a higher water content. The
78 decontamination of dried powders is difficult, and the difficulty correlates with the increasing
79 resistance of spores and vegetative forms to low water activity levels (Fine and Gervais,
80 2005b; Laroche et al., 2005; Laroche and Gervais, 2003; Tiburski et al., 2014).

81 Many authors have already shown that the low water activity or low water content in spores
82 could explain their high heat resistance and that water diffusion from or out of the spore core
83 is slowed (Knudsen et al., 2016; Loison et al., 2013; Nguyen Thi Minh et al., 2010b). In a
84 previous work, *Bacillus subtilis* spores have been reported to have a low water permeability
85 and suggested that the coat structure may be necessary for maintaining this low a_w (Knudsen
86 et al., 2016). In another manuscript, using a molecular rotor, the high viscosity of the internal
87 membrane of this spore was demonstrated and correlated this property with its barrier
88 properties (Loison et al., 2013).

89 In previous studies on spore thermal inactivation, *Bacillus subtilis* spores were shown to have
90 D-values which fell in the same range of pathogen spores like *B. cereus* and *B. anthracis*
91 (Montville et al., 2005). Nevertheless, large interstrain variations in spore heat resistance has
92 been regularly reported. For example, the presence of a specific operon (SPOVA2mob not
93 present in strain 168 used in this work) leads to increase the D value more than 100 times
94 (Krawczyk et al., 2017).

95 *B. subtilis* is also used as a surrogate for some pathogenic spores (Hu and Gurtler, 2017).
96 Moreover *B. subtilis* itself can induce food spoilage incidents particularly on bread
97 (Thompson et al., 1993).

98 Recently Tros et al. (2017) have shown that a large fraction of “slow” intracellular water
99 linked to proteins exists in the *Bacillus subtilis* spores and in a previous work (Tiburski et al.,
100 2014), the role of water in the heat resistance of dried *Bacillus subtilis* spores was also
101 investigated using Differential Scanning Calorimetry. The spore temperature was linearly
102 increased in pans with different pressure resistances. The results demonstrated that even in
103 dry external conditions, a water fraction stayed embedded in the spore core and that
104 maintenance of this core water content during heating dramatically increased spore
105 destruction. Infrared analysis of treated spores showed that the heat destruction was highly
106 related to protein denaturation and dipicolinic acid (DPA) release, which were maximized in
107 wet conditions.

108 This work was intended to apply these results to optimizing dried spore decontamination
109 based on preserving the initial water content of the spore core. The method used to prevent
110 water evaporation was to pressurize a hermetic container and consequently change the water
111 vapor equilibrium.

This study investigated the inactivation of dried *Bacillus subtilis* spores equilibrated at different water activities by a treatment using high temperature and absolute pressure levels from 0.1 to 0.6 MPa. *Bacillus subtilis* spores were already used as surrogates for pathogenic spores (Hu and Gurtler, 2017). The applied pressure prevented the water in the samples from boiling, thus allowing its initial water content to be maintained. The effect of time/temperature treatments under different pressure levels and for different initial water activities on final spore viability were compared.

2. Material and Methods

2.1 Bacterial strain and growth conditions

The reference strain *B. subtilis* 168 from the BGSC (Bacillus Genetic Stock Center, Department of Biochemistry, The Ohio State University, USA) was used in this study. *B. subtilis* sporulation was induced in a complex medium in a reactor at 37°C, pH 8.0, with an air flow of 4 L/min and with agitation (450 rpm) (Nguyen Thi Minh et al., 2008). Sporulation was estimated by plating on Bromocresol Purple agar (Dextrose Tryptone agar, BIODIA Diagnostics, Beauvais, France) before and after a heat treatment at 80°C for 10 min. When more than 95% of the population in the sample resisted this heat treatment, the spore suspension was harvested (3–5 days). The spore suspension was washed with sterile distilled water four times and was then spray-dried (Mini spray dryer B-290, Buchi, France) and stored in powder form in sterile receptacles at 4°C until use. The final microbial concentration of the powder was approximately 10^{11} CFU/g.

2.2 Equilibration of spores at different water activities in gaseous and liquid media

Dried spores with water activity values of 0.10, 0.20, 0.30, 0.40, 0.50 and 0.70 were obtained by placing the spores (± 500 mg) inside 1 L air-tight plastic boxes containing the water/glycerol solutions presented in Table 1 (the solution occupied 1/10 of the box volume) until equilibrium was reached (at least one week). Water activity determinations of the osmotic solutions and spore powders were performed in triplicate at 25°C using a Decagon-AQUALAB CX-2 water activity meter (Meter, WA, USA).

For liquid media, 10 mg of spray-dried spores previously equilibrated at different water activity levels were placed directly in water/glycerol solutions at concentrations predicted by the Norrish Equation to correspond to a_w levels of 0.10 to 0.70 (Norrish, 1966). The experimental a_w values were also checked with the osmometer previously described.

2.3 Treatment of dried spores in gaseous or liquid medium at high temperature and up to a pressure level of 0.6 MPa

The reactor consisted of a small hermetic stainless-steel cylinder (1 cm³ of volume) that could be pressurized with gas. Nitrogen was used to pressurize the reactor, after which the reactor's valve was closed to maintain the applied pressure.

2.3.1 Gaseous medium

In closed systems containing water, increasing pressure prevents water vaporization. Therefore, if the pressure in the reactor is lower than the pressure of water vaporization (P_{vap}) at the considered temperature, water vaporization from the spores will occur until P_{vap} is reached. On the other hand, as soon as the pressure exceeds P_{vap} , no vaporization of the water present in the spores will occur. The final pressure inside the reactor is determined by the input controlled pressure of nitrogen and two modifying factors: i) the pressure increase resulting from increased temperature and ii) the pressure increase resulting from water evaporation from the sample. These factors have been considered to fit the evolution of the

water vaporization in the reactor as a function of the total pressure evolution for different initial nitrogen pressures.

Figure 1 models the increasing pressure in the vessel for each initial nitrogen pressure level as a function of the increasing temperature and the water vaporization temperature evolution curve. As soon as the temperature/pressure curve crosses the temperature vaporization curve (which occurs for 0.1 MPa, 0.2 MPa and 0.3 MPa at 107, 130 and 145°C, respectively) any liquid water present in the vessel will evaporate from this intersection point until the end of the heat treatment. For the 3 higher initial pressure levels (i.e., 0.4, 0.5 and 0.6 MPa), the 3 curves did not cross the water saturation curve before a temperature of 150°C was reached in the experiment (see Figure 1), thus one can assume that there was no evaporation of any liquid water in the vessel from an initial pressure of 0.4 MPa.

For most experiments, 10 mg of dried spores were placed in aluminum foil that was folded and placed inside the reactor. The reactor was then pressurized between 0.2 and 0.6 MPa, closed and heated in an oil bath at 150°C. When atmospheric pressure was used, the valve was left open to allow water evaporation. After the treatment, the reactors were cooled in an ice bath, then the valve was opened to release the pressure.

2.3.2 Liquid medium

For one set of experiments, performed at 0.6 MPa initial pressure and 150°C, the reactor described previously was filled with water-glycerol suspensions at the same five a_w values previously examined in gaseous medium, and 10 mg of spores were added immediately before heat treatment. Under these thermodynamic conditions, there was no vaporization of the different water/glycerol solutions or of pure water.

For liquid and gaseous treatment, it was verified that such a pressure level (0.6 MPa) does not alter spore viability. In fact pressure levels greater than 200 MPa are necessary to slightly inactivate spores with gas medium (Colas de la Noue et al., 2012) and 100 MPa in liquid medium (Nguyen Thi Minh et al., 2010a).

186

2.4 Spore viability

Heat-treated dried spores and spores in liquid medium were serially diluted with physiological saline solution (0.9% NaCl w/v), and 100 µl of each dilution was seeded in triplicate onto BCP agar (Dextrose Tryptone Agar, Biokar Diagnostics, Beauvais, France) plates. Colonies were counted after 24 hours of incubation at 37°C. Control samples contained untreated spores. All experiments were performed at least in triplicate. Spore inactivation was expressed using a logarithmic reduction factor $\log (N_0 - N)/N_0$, with N_0 representing the number of developing spores before treatment and N representing the number of developing spores after treatment.

2.5 Statistical analysis

On each viability point (at least 3 independent measurements) standard deviation was calculated and plotted as error bars on each figure. In order to confirm the effect of pressure level, an ANOVA and Tukey's honestly significant difference test (if $p < 0.05$) were performed to determinate whether significant differences existed among treatment at different pressure. Analyses were performed using the R software, version 3.1.2.

3. Results

3.1 Effect of temperature and pressure on the viability of dried spores in gaseous medium

Initial pressures between 0.2 and 0.6 MPa were used to study the effect of pressure combined with heat in *Bacillus* spore inactivation. Spores were equilibrated at a water activity of 0.50, which is a typical a_w for dried foods such as flours, cereals and spices (Lang et al., 2017). The heat treatment (150°C, 120 s) was chosen after preliminary tests that showed these conditions to be discriminative on the ratio of spore viability to mortality.

In Figure 2, we show that heat treatment performed with no additional pressure, i.e., 150°C at atmospheric pressure, resulted in a slight destruction level of 0.3 log, which is not significantly different from the control sample maintained at 25°C and which certainly corresponds to the inactivation of pre-germinating or germinating spores.

Thus, this treatment did not inactivate any spores, which illustrates their significant thermoresistance. The results presented in Figure 2 also clearly show that when the applied pressure was increased, spore inactivation also increased. Below 0.3 MPa, the inactivation level was found to be less than 3 log, whereas above 0.3 MPa, the inactivation suddenly increased from 4 log CFU for 0.4 MPa to 5 log CFU for 0.6 MPa.

An ANOVA performed on these results confirmed that the inactivation levels corresponding to the two pressure level groups (0.2 and 0.3 MPa versus 0.4, 0.5 and 0.6 MPa) were significantly different ($p < 0.05$).

To further probe the previous results, the kinetics of spore inactivation was studied at a_w 0.5 and 150°C. The pressure used was 0.6 MPa, which corresponded to higher spore destruction in Figure 2. The inactivation of spores with time is shown in Figure 3. Spores heated at atmospheric pressure exhibited only a slight inactivation of approximately 0.5 log after 240 s, whereas spores subjected to 0.6 MPa pressure exhibited an inactivation of 5 log between 60 s and 120 s of treatment.

Data from Figures 2 and 3 show that pressure significantly affects the destruction of *B. subtilis* spores undergoing a 150°C heat treatment at an a_w of 0.5. This pressure effect exists at all pressure levels but becomes more significant with a gap of 1 to 2 log destruction for initial pressure levels greater than 0.4 MPa.

3.2 Effect of water activity on the viability of dried spores at different times for a 150°C/0.6 MPa treatment

Dried food products and food powders normally have a water activity ranging from 0.20 to 0.50, and in exceptional cases, their a_w may reach 0.70, thus it is important to investigate the effect of this variable on the proposed treatment.

3.2.1 In gaseous medium

The influence of water activity on spore inactivation is presented in Figure 4. These results clearly show that the water activity, and consequently the water content of the spores, plays a fundamental role in spore resistance. Spores equilibrated at an a_w of 0.10 exhibited an inactivation of 5 log after 560 s, whereas spores at an a_w of 0.20 exhibited an inactivation of 8 log after the same amount of time. When water activity increased, spore inactivation also increased, as shown by the decrease in logarithmic reduction time (D-values) presented in Table 2. Indeed, the D value is four times greater for an a_w value of 0.10 than for an a_w value of 0.7. Moreover, we found that the same increase in water activity level does not result in a proportional increase in spore inactivation. For example, there is only a small difference between spore inactivation at a_w 0.30 and 0.50, but further increase to an a_w of 0.70 resulted in complete inactivation after only 240 s.

It is important to emphasize that, with a pressure of 0.6 MPa, the treatment times are greatly reduced, and so, even for water activity levels as low as 0.10 and 0.20, it is possible to reduce the spore population by a factor of 10 to 5 in approximately 9 or 6 minutes, respectively.

3.2.2 In liquid medium

To build on the previous results, the protective effect of a_w on spore thermal resistance was analyzed in a liquid aqueous medium in the same experimental chamber used for spore inactivation in a gaseous medium. Addition of glycerol in the treatment liquid medium was used to control the a_w of the medium and therefore may have protected spores from heat. However, osmolytes sometimes also have direct toxic effects on spores and interfere with the pure osmotic effect (Mazas et al., 1999). Nevertheless, spores are known for their low permeability to many toxic chemicals, in particular, those that can damage the spore DNA located in the central spore core (Knudsen et al., 2016). Moreover, glycerol, which was used in our experiment, is a compatible solute with very few interactions with other molecules, thus it is an almost ideal solute. Therefore, its action may be mainly related to the a_w decrease.

The results shown in Figure 5 demonstrate that the water activity of the treatment medium had a major effect on spore heat inactivation. When heated in higher water activities, spores were more sensitive to heat. Spores treated at a_w 0.10 or 0.20 presented a 2 log reduction after 240 s, whereas spores at a_w 0.30, 0.40 and 0.50 presented a 6 log inactivation after only 180 s.

Spore inactivation began at different treatment times depending on the a_w of the medium and later than we observed for spores treated in dried form (see Figure 3). This result can be explained by the heating kinetics inside the reactor. Indeed, previous experiments (data not shown) have allowed us to verify that the temperature inside the reactor reached 99% of the oil bath temperature in less than 10 s when air/nitrogen was the transmitting medium;

however, more than 100 s were needed for pure glycerol and 60 s for pure water to reach the same 99% temperature response. Figure 5 shows that inactivation at an a_w of 0.70 was noticeable between 60 s and 90 s, whereas for lower a_w levels, corresponding to lower spore water content, inactivation started only after 120 s of treatment.

Comparing these results with those obtained from dried spores in gaseous conditions (see Figure 3), it can easily be noted that for the same water activity, the final spore inactivation was quite similar.

4. Discussion

Steam treatment is in extensive use today for microbial decontamination of dry foods, but its use presents problems since it results in water condensation on the product that must be removed to avoid subsequent mold growth and spoilage. On the other hand, thermal processes such as microwaves (Kim et al., 2009), infrared (Eliasson et al., 2014) and High Temperature Short Time treatment (Fine and Gervais, 2005a) necessarily involve supplementary drying of the product. Moreover, alternative treatments available for use in aqueous media, such as high pressure and pulsed electric fields, are not efficient for the decontamination of dried foods (Espinasse et al., 2008). Therefore, new procedures for decontaminating dried foods that can destroy bacterial spores are needed.

This study investigates the inactivation of dried *Bacillus subtilis* spores equilibrated at different water activities by a treatment using high temperature and absolute pressure levels of nitrogen from 0.2 to 0.6 MPa. In this experiment, the applied pressure prevented water evaporation from the treated samples, allowing the initial water content of the spores, which is mainly situated in the spore core, to be maintained.

300 Inactivation varies from less than 1 log at atmospheric pressure to more than 5 log at 0.6 MPa.
301 Between the initial pressure level of 0.3 MPa at 25°C (which generates 0.43 MPa at 150°C)
302 and the initial pressure 0.4 MPa at 25°C (which generates 0.57 MPa at 150°C), there is a
303 sudden increase in inactivation, in contrast to the slower increase observed at higher
304 pressures. Based on these results, it can be concluded that when the pressure generated in the
305 vessel exceeded the pressure corresponding to the water vaporization temperature of 150°C
306 (0.47 MPa, see Figure 1) the spore inactivation significantly increased, as shown in Figure 2.

307 From this result, it can be proposed that i) maintaining water in the endospore favors bacteria
308 inactivation and that ii) spore drying, which is initiated for pressure levels lower than 0.47
309 MPa, favors bacteria survival at 150°C. Moreover, the drying time and thus the final drying
310 level decreased with increasing pressure until no further drying occurred, when the internal
311 pressure reached the 0.47 MPa value, which corresponded to a nitrogen pressure between 0.3
312 and 0.4 MPa (see Figure 1).

313 One can propose a role of water in spore inactivation in agreement with numerous works
314 (Coleman et al., 2007; Sochava, 1997; Subramanian et al., 2006) in which the internal
315 endospore macromolecules, mainly composed of DNA, DPA, and proteins (the last being the
316 most delicate), are preferentially heat-damaged when mobile and unfolded in a water medium.
317 Moreover, a recent work (Tros et al., 2017) has observed a large fraction of “normal”
318 intracellular water in the spore core of *B. subtilis*.

319 Even for very low water activity levels, noticeable spore inactivation was obtained in liquid as
320 well as in gaseous medium. These results agree with previous work (Tiburski et al., 2014),
321 which has shown a residual high water content in the core even for low a_w -equilibrated
322 spores. This phenomenon was attributed to specific high barrier properties of the internal

spore membrane slowing down the thermodynamic balance. Thus, it could be assumed that some water remaining in the core caused macromolecule denaturation.

Inactivating one or more key spore proteins is believed to be important in spore heat killing (Coleman et al., 2010). When heat surpasses a threshold level, especially in aqueous media, changes in the structure of vital growth proteins become irreversible, causing spore inactivation (Grinshpun et al., 2010). It is also known that in low moisture states the temperature of protein denaturation increases (Sochava, 1997), making inactivation of these key proteins more difficult. In this context, the high spore heat resistance at low water content can be explained by the low mobility and changes in the conformations of proteins and enzymes. Indeed, most spore proteins are rotationally immobilized, which may contribute to heat resistance by preventing heat-denatured proteins from aggregating irreversibly (Sunde et al., 2009). Consequently, when the original water content of the spore is maintained, protein damage is increased during heat treatment, and such damage is certainly relevant to the differences in spore inactivation found in our experiments. This hypothesis has been directly verified through FTIR measurement in a previous work (Tiburski et al., 2014). Nevertheless, the present results were obtained on *B. subtilis* strain 168 and it is possible that at the pressure and temperature levels used the spore inactivation mechanisms could vary for other bacterial spores.

Conclusion

The model system developed in this work can be extended to treating low water content foods. This system is based on the application of temperature under pressure, which prevents any water evaporation from bacterial spores embedded in the food product. This model has the advantage of not adding moisture to the product, and it results in high spore destruction rates (approximately 5 log). The efficiency of this process is related to the high water

permeability barrier of the internal membrane of the spore, which has been shown to allow maintenance of a core water content greater than the outer membrane long after drying. Thus, this new process is particularly efficient for dried bacteria spore destruction because of their high residual core water content. Further studies must first verify this effect on other sporulated strains and especially pathogens. In a second phase, we must integrate the impact of such pressure/temperature/time treatments on the organoleptic and nutritional properties of different dried food products. Indeed, optimal treatments should conserve the functional properties of food in addition to assuring maximal bacteria spore inactivation. Following this work, this process could then be tested to treat dried food products such as spices. The normal a_w of dried food ranges between 0.30 and 0.50, and in this work, we have shown that spore inactivation higher than 5 log can be expected from a 3 minute treatment at these water activities. Therefore, if the products already possess a similar a_w , no previous equilibration will be needed; however, if the product to be processed has a very low a_w , an equilibration step would be necessary. In this case, the dried food could be stored in an atmosphere with a relative humidity of 50% for a few weeks prior to treatment. Certainly, different conditions of time, temperature and pressure adapted to each product will need to be tested to optimize this new decontamination procedure.

The next step would be to develop a scaled-up version of the system to perform tests in real food products to evaluate both microbiological decontamination and changes in quality parameters.

Acknowledgments

This work is supported by the Conseil Régional de Bourgogne through the plan d'actions régional pour l'innovation (PARI) and the European Union through the PO FEDER-FSE

371 Bourgogne 2014/2020 programs. The PhD of Julia Hauck Tiburski was funded by the
372 Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES-Brazil).

373

374 References

375 Andersson, A., Rönner, U., Granum, P.E., 1995. What problems does the food industry have with the
376 spore-forming pathogens *Bacillus cereus* and *Clostridium perfringens* ? Int. J. Food Microbiol. 28(2),
377 145-155.

378 Colas de la Noue, A., Espinasse, V., Perrier-Cornet, J.M., Gervais, P., 2012. High gas pressure: an
379 innovative method for the inactivation of dried bacterial spores. Biotechnol. Bioeng. 109(8), 1996-
380 2004.

381 Coleman, W.H., Chen, D., Li, Y.Q., Cowan, A.E., Setlow, P., 2007. How moist heat kills spores of
382 *Bacillus subtilis*. J. Bacteriol. 189(23), 8458–8466.

383 Coleman, W.H., Zhang, P., Li, Y.Q., Setlow, P., 2010. Mechanism of killing of spores of *Bacillus cereus*
384 and *Bacillus megaterium* by wet heat. Lett. Appl. Microbiol. 50(5), 507-514.

385 Cuq, B., Rondet, E., Abecassis, J., 2011. Food powders engineering, between knowhow and science:
386 Constraints, stakes and opportunities. Powder Technol. 208(2), 244-251.

387 Doyle, C.J., Gleeson, D., Jordan, K., Beresford, T.P., Ross, R.P., Fitzgerald, G.F., Cotter, P.D., 2015.
388 Anaerobic sporeformers and their significance with respect to milk and dairy products. Int. J. Food
389 Microbiol. 197(1), 77-87.

390 Eliasson, L., Libander, P., Lövenklev, M., Isaksson, S., Ahrné, L., 2014. Infrared decontamination of
391 oregano: effects on *Bacillus cereus* spores, water activity, color, and volatile compounds. J. Food Sci.
392 79(12), 2447-2455.

393 Espinasse, V., Perrier-Cornet, J.M., Marecat, A., Gervais, P., 2008. High-Pressure inactivation of dried
394 microorganisms. Biotechnol. Bioeng. 99(1), 180-188.

395 Fine, F., Gervais, P., 2005a. A new high temperature short time process for microbial
 396 decontamination of seeds and food powders. *Powder Technol.* 157(1-3), 108-113.

397 Fine, F., Gervais, P., 2005b. Thermal destruction of dried vegetative yeast cells and dried bacterial
 398 spores in a convective hot air flow: strong influence of initial water activity. *Environ. Microbiol.* 7(1),
 399 40-46.

400 Ghisleni, D., Braga, M., Kikuchi, I., Braşoveanu, M., Nemţanu, M., Dua, K., Pinto, T., 2016. The
 401 microbial quality aspects and decontamination approaches for the herbal medicinal plants and
 402 products: an in-depth review. *Curr. Pharm. Des.* 22, 4264-4287.

403 Glasset, B., Herbin, S., Guillier, L., Cadel-Six, S., Vignaud, M.L., Grout, J., Pairaud, S., Michel, V.,
 404 Hennekinne, J.A., Ramarao, N., Brisabois, A., 2016. *Bacillus cereus*-induced food-borne outbreaks in
 405 France, 2007 to 2014: epidemiology and genetic characterisation. *Euro Surveill.* 21(48), 30413.

406 Grasso, E.M., Stam, C.N., Anderson, N.M., Krishnamurthy, K., 2014. Heat and steam treatments, in:
 407 Gurtler, J.B., Doyle, M.P., Kornacki, J.L. (Eds.), *The microbiological safety of low water activity foods*
 408 *and spices*. Springer, New York, pp. 403-424.

409 Grinshpun, S.A., Adhikari, A., Li, C., Reponen, T., Yermakov, M., Schoenitz, M., Dreizin, E., Trunov, M.,
 410 Mohan, S., 2010. Thermal inactivation of airborne viable *Bacillus subtilis* spores by short-term
 411 exposure in axially heated air flow. *J. Aerosol Sci.* 41(4), 352-363.

412 Hu, M., Gurtler, J.B., 2017. Selection of surrogate bacteria for use in food safety challenge studies: a
 413 review. *J. Food Prot.* 80(9), 1506-1536.

414 Kim, S.Y., Shin, S.J., Song, C.H., Jo, E.K., Kim, H.J., Park, J.K., 2009. Destruction of *Bacillus licheniformis*
 415 spores by microwave irradiation. *J. Appl. Microbiol.* 106(3), 877-885.

416 Knudsen, S.M., Cermak, N., Feijó Delgado, F., Setlow, B., Setlow, P., Manalis, S.R., 2016. Water and
 417 small-molecule permeation of dormant *Bacillus subtilis* spores. *J. Bacteriol.* 198(1), 168-177.

418 Krawczyk, A.O., de Jong, A., Omony, J., Holsappel, S., Wells-Bennik, M.H.J., Kuipers, O.P., Eijlander,
 419 R.T., 2017. Spore heat-activation requirements and germination responses correlate with sequences

420 of germinant receptors and with the presence of a specific *spoVA*^{2mob} operon in food-borne strains of
 421 *Bacillus subtilis*. Appl. Environ. Microbiol. 83(7), e03122-16.
 422 Lang, E., Chemlal, L., Molin, P., Guyot, S., Alvarez-Martin, P., Perrier-Cornet, J.M., Dantigny, P.,
 423 Gervais, P., 2017. Modeling the heat inactivation of foodborne pathogens in milk powder: High
 424 relevance of the substrate water activity. Food Res. Int. 99(1), 577-585.
 425 Laroche, C., Gervais, P., 2003. Unexpected thermal destruction of dried, glass bead-immobilized
 426 microorganisms as a function of water activity. Appl. Environ. Microbiol. 69(5), 3015-3019.
 427 Laroche, C., Fine, F., Gervais, P., 2005. Water activity affects heat resistance of microorganisms in
 428 food powders. Int. J. Food Microbiol. 97(3), 307-315.
 429 Logan, N.A., 2012. *Bacillus* and relatives in foodborne illness. J. Appl. Microbiol. 112(3), 417-429.
 430 Loison, P., Hosny, N.A., Gervais, P., Champion, D., Kuimova, M.K., Perrier-Cornet, J.M., 2013. Direct
 431 investigation of viscosity of an atypical inner membrane of *Bacillus* spores: a molecular rotor/FLIM
 432 study. Biochim. Biophys. Acta 1828(11), 2436-2443.
 433 Mazas, M., Martínez, S., López, M., Alvarez, A.B., Martin, R., 1999. Thermal inactivation of *Bacillus*
 434 *cereus* spores affected by the solutes used to control water activity of the heating medium. Int. J.
 435 Food Microbiol. 53(1), 61-67.
 436 Montville, T.J., Dengrove, R., De Siano, T., Bonnet, M., Schaffner, D.W., 2005. Thermal resistance of
 437 spores from virulent strains of *Bacillus anthracis* and potential surrogates. J. Food Prot. 68(11), 2362-
 438 2366.
 439 Nguyen Thi Minh, H., Perrier-Cornet, J.M., Gervais, P., 2008. Effect of the osmotic conditions during
 440 sporulation on the subsequent resistance of bacterial spores. Appl. Microbiol. Biotechnol. 80, 107-
 441 114.
 442 Nguyen Thi Minh, H., Dantigny, P., Perrier-Cornet, J.M., Gervais, P., 2010a. Germination and
 443 inactivation of *Bacillus subtilis* spores induced by moderate hydrostatic pressure. Biotechnol. Bioeng.
 444 107(5), 876-883.

445 Nguyen Thi Minh, H., Perrier-Cornet, J.M., Gervais, P., Champion, D., 2010b. New Insight into the
 446 thermal properties and the biological behaviour of the bacterial spores. *Food Biophys.* 5(3), 238-246.
 447 Norrish, R.S., 1966. An equation for the activity coefficients and equilibrium relative humidities of
 448 water in confectionery syrups. *J. Food Technol.* 1, 25-39.
 449 Silva, M.P., Pereira, C.A., Junqueira, J.C., Jorge, A.O.C., 2013. Methods of destroying bacterial spores,
 450 in: Méndez-Vilas, A. (Ed.), *Microbial pathogens and strategies for combating them: science,*
 451 *technology and education.* Formatex Research Center, Spain, pp. 490-496.
 452 Sochava, I.V., 1997. Heat capacity and thermodynamic characteristics of denaturation and glass
 453 transition of hydrated and anhydrous proteins. *Biophys. Chem.* 69(1), 31-41.
 454 Subramanian, A., Ahn, J., Balasubramaniam, V.M., Rodriguez-Saona, L., 2006. Determination of spore
 455 inactivation during thermal and pressure-assisted thermal processing using FT-IR Spectroscopy. *J.*
 456 *Agric. Food Chem.* 54(26), 10300-10306.
 457 Sunde, E.P., Setlow, P., Hederstedt, L., Halle, B., 2009. The physical state of water in bacterial spores.
 458 *Proc. Natl. Acad. Sci. USA* 106(46), 19334–19339.
 459 Tauveron, G., Slomianny, C., Henry, C., Faille, C., 2006. Variability among *Bacillus cereus* strains in
 460 spore surface properties and influence on their ability to contaminate food surface equipment. *Int. J.*
 461 *Food Microbiol.* 110(3), 254-262.
 462 Thompson, J.M., Dodd, C.E.R., Waites, W.M., 1993. Spoilage of bread by *Bacillus*. *Int. Biodeterior.*
 463 *Biodegrad.* 32(1), 55-66.
 464 Tiburski, J.H., Rosenthal, A., Guyot, S., Perrier-Cornet, J.M., Gervais, P., 2014. Water distribution in
 465 bacterial spores: A key factor in heat resistance. *Food Biophys.* 9(1), 10-19.
 466 Tros, M., Zheng, L., Hunger, J., Bonn, M., Bonn, D., Smits, G.J., Woutersen, S., 2017. Picosecond
 467 orientational dynamics of water in living cells. *Nat. Commun.* 8(1), 904.
 468 Van Doren, J.M., Neil, K.P., Parish, M., Gieraltowski, L., Gould, L.H., Gombas, K.L., 2013. Foodborne
 469 illness outbreaks from microbial contaminants in spices, 1973–2010. *Food Microbiol.* 36(2), 456-464.

Figure 1: Calculated variations of internal pressure during isochoric heating of pressurization devices (internal volume of 1.7 ml) loaded with a 10 mg spores sample initially equilibrated at a_w 0.5 and at initial pressure ranging from 0.1 MPa to 0.6 MPa, in comparison with water boiling temperature of a model saline solution initially at a_w 0.5. (NIST Chemistry Webbook has been used for water data calculation).

Figure 2: Thermal inactivation of *B. subtilis* 168 dried spores initially equilibrated at a water activity of 0.50 treated at 150°C / 120s in a gaseous medium at different initial pressure levels. (Error bars are SD, Mean comparison using Tukey's test: same letters indicate no significant difference between treatments $P>0.95$)


Figure 3: Kinetics of *B. subtilis* 168 dried spores' inactivation. The spores have been dried and equilibrated at a water activity of 0.50 and treated at 150°C in a gaseous medium initially at atmospheric pressure (empty triangle) or at 0.6 MPa (dot). Error bars are SD of at least 3 independent measures.


Figure 4: Inactivation of *B. subtilis* 168 spores equilibrated at different water activities (0.10, 0.20, 0.30, 0.50, and 0.70) after a treatment at 150°C and 0.6 MPa. Error bars are SD of at least 3 independent measures


Figure 5: Inactivation of *B. subtilis* 168 spores in water/glycerol binary solutions at different water activities after a treatment at 150°C and 0.6 MPa. Error bars are SD of at least 3 independent measures.






Table 1: Glycerol concentration (from Norrish equation) used to equilibrate the water activity in water/glycerol binary solutions.


Table 2: Decimal reduction times (in minutes) for *B. subtilis* 168 dried spores according to the initial water activity of gaseous environment for a treatment at 150°C and a pressure of 0.6 MPa











a_w	Glycerol (g/100g solution)
0.10	96
0.20	92
0.30	88
0.40	84
0.50	77
0.60	72
0.70	64

a_w	D (min)
0.10	1.7
0.20	1.0
0.30	0.7
0.50	0.8
0.70	0.5