

Does the Motor Cortex Want the Full Story? The Influence of Sentence Context on Corticospinal Excitability in Action Language Processing

W. Dupont, C. Papaxanthis, Florent Lebon, Carol Madden-Lombardi

▶ To cite this version:

W. Dupont, C. Papaxanthis, Florent Lebon, Carol Madden-Lombardi. Does the Motor Cortex Want the Full Story? The Influence of Sentence Context on Corticospinal Excitability in Action Language Processing. Neuroscience, 2022, 506, pp.58-67. 10.1016/j.neuroscience.2022.10.022. hal-04185286

HAL Id: hal-04185286 https://u-bourgogne.hal.science/hal-04185286v1

Submitted on 10 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1	Does the motor cortex want the full story? The influence of sentence context
2	on corticospinal excitability in action language processing
3	
4	
5	Dupont W 1*, Papaxanthis C 1, Lebon F 1#, Madden-Lombardi C 12#
6	¹ INSERM UMR1093-CAPS, Université Bourgogne Franche-Comté, UFR des
7	Sciences du Sport, F-21000, Dijon. ² Centre National de la Recherche Scientifique
8	(CNRS), France
9	
10	
11	# F.L. and C. M-L. equally contributed to this work
12	* Corresponding author: william.dupont@u-bourgogne.fr
13	

14 Highlights

- We studied the effects of action words and sentence context on motor activity.
- Corticospinal excitability is modulated by the presence of manual action verbs.
- Context helps eliminate residual motor activation when no manual action is present.

Abstract

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

The reading of action verbs has been shown to activate motor areas, whereby sentence context may serve to either globally strengthen this activation or to selectively sharpen it. To investigate this issue, we manipulated the presence of manual actions and sentence context, assessing the level of corticospinal excitability by means of transcranial magnetic stimulation. We hypothesized that context would serve to sharpen the neural representation of the described actions in the motor cortex, reflected in context-specific modulation of corticospinal excitability. Participants silently read manual action verbs and non-manual verbs, preceded by a full sentence (rich context) or not (minimal context). Transcranial magnetic stimulation pulses were delivered at rest or shortly after verb presentation. The coil was positioned over the cortical representation of the right first dorsal interosseous (pointer finger). We observed a general increase of corticospinal excitability while reading both manual action and non-manual verbs in minimal context, whereas the modulation was action-specific in rich context: corticospinal excitability increased while reading manual verbs, but did not differ from baseline for non-manual verbs. These findings suggest that sentence context sharpens motor representations, activating the motor cortex when relevant and eliminating any residual motor activation when no action is present.

38 Keywords: Action language, motor representation, transcranial magnetic stimulation.

Introduction

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

Over the past couple of decades, researchers have established a link between language processes and perceptual-motor processes (Barsalou, 1999, 2008; Barsalou et al., 2003; Gallese and Lakoff, 2005; Decety and Grèzes, 2006; Zwaan and Taylor, 2006; Fischer and Zwaan, 2008; Jeannerod, 2008; Glenberg and Gallese, 2012). More precisely, these authors suggest that the information described by language would lead to activation in the brain areas that are concerned with the concept itself; if the concept concerns sight, this will lead to activation in the visual areas. The same applies to action language, such that semantic information referring to an action would engage the sensory motor system. While there exist inconsistencies in the literature suggesting that this distributed activation may be epiphenomenal at most (Rogers et al., 2006; Patterson et al., 2007; Mahon and Caramazza, 2008; Lambon Ralph et al., 2009, 2010; Pobric et al., 2010; Mahon, 2015), many researchers would postulate that such activation contributes to language comprehension.

To better understand this link, researchers have probed the involvement of the motor system during action language processing using various methodologies, such as functional Magnetic Resonance Imaging (fMRI) (Hauk et al., 2004; Pulvermüller et al., 2005; Aziz-Zadeh et al., 2006; Van Dam et al., 2010; although for critical views see: de Zubicaray et al., 2013; Wurm and Caramazza, 2019). electroencephalography (EEG) (Pulvermüller et al., 2001; Kellenbach et al., 2002; Xu et al., 2016), behavioral measures (Pulvermüller et al., 2001; Rabahi et al., 2012, 2013; Andres et al., 2015; Klepp et al., 2019) and Transcranial Magnetic Stimulation (TMS) (Papeo et al., 2009, 2015; Labruna et al., 2011; Scorolli et al., 2012; Innocenti et al., 2014; Franklin et al., 2020; see review: Papeo et al. 2013). Specifically, Hauk et al. (2004) using fMRI reported that silent reading of action words leads to somatotopic activation of the effector-related representation within the primary motor cortex. Papeo et al. (2009) used TMS to observe an increase in MEP amplitude for hand action verbs vs. non-action verbs 500 ms after verb presentation. These cortical activations could explain behavioral modulation during language processing, such as improvement in squat jump (Rabahi et al., 2012, 2013), and facilitation of motor response time (Andres et al., 2015; Klepp et al., 2019).

Such behavioral facilitations are present only if the linguistic content is compatible with the movement performed (Gentilucci et al., 2000; Taylor and Zwaan, 2008; Dalla Volta et al., 2009; Glenberg and Gallese, 2012; Zwaan et al., 2012; Van Dam and Desai, 2017). Furthermore, some studies report decreases in motor activation for action verbs compared to non-action verbs at very early latencies (TMS before end of verb: Buccino et al., 2005), or fail to observe activation of the motor cortex (lexical decision tasks: Perani et al., 1999; Longe et al., 2007; motion word comprehension: Grossman et al., 2002; passive reading: Gianelli et al., 2020). While some of the discrepancies in these findings may be attributable to the timing of these studies (see early interference explanation proposed by Buccino et al., 2005), such results generally undermine the idea that action verbs systematically/automatically activate the motor cortex (see also Rüschemeyer et al., 2007; Raposo et al., 2009). Instead, this activation seems to be, as Kemmerer (2015, page 4) states, "sensitive to attentional and situational factors that we are only beginning to understand."

In our view, context is one of these factors that plays an important role in language processing, allowing a more effective and precise understanding of the action described in the sentence. Indeed, studies using fMRI and behavioral measures demonstrate how context can modulate the degree of motor activation and

performance (Raposo et al., 2009; Van Dam et al., 2010; van Dam et al., 2010; Zwaan et al., 2010; Papeo et al., 2012; Van Dam et al., 2012; van Dam et al., 2012; Gilead et al., 2013; Van Dam et al., 2014; Kemmerer, 2015; Beauprez et al., 2018). For instance, Raposo and colleagues (2009) observed activation of motor areas for action verbs in isolation (e.g., *kick*) and in literal phrases (e.g., *kick the ball*), but not in idiomatic phrases (e.g., *kick the bucket*). Likewise, Van Dam et colleagues (2010) observed greater activation in motor-related brain areas for highly specific action verbs (to wave, to wipe) than for less specific action verbs (to greet, to clean).

Given these results, it is possible that specifying actions by increasing the sentence context may lead to greater motor cortex activation in general. Indeed, Wurm and Caramazza (2019) report a stronger decoding of action scenes relative to short sentences in frontoparietal representations, which they attribute to a difference in richness of detail. Although motor representations can be triggered by the reading or perception of an action, these researchers suggest that action reading leads to a less specific and robust motor representation. Following this logic, increasing sentence context (specifying the action) could increase motor cortex activation in general, thereby "strengthening" motor representations.

Another possibility is that greater sentence context leads to a "sharpening" of motor representations, making them more precise yet not always stronger. This idea is consistent with studies in which the context surrounding action verbs selectively engages motor representations. For instance, Taylor and Zwaan (2008) observed facilitation for rotating a knob when the direction of rotation (clockwise) matched rather than mismatched the direction described in concurrent sentences (*The gardener noticed that the water was still running. He approached the faucet which he turned off quickly*). Interestingly, the motor facilitation was observed at the moment

of processing the verb, and reappeared at the post-verbal adverb, only if that adverb qualified the action ("slowly" or "quickly"), and not if the adverb instead shifted focus to the agent (e.g., "obediently" or "eagerly"). This suggests that context can selectively activate motor representations, and furthermore, that these motor representations are temporally pinned to the verb, or any word that draws the contextual focus to the action itself.

In the current study, we investigated whether context leads to a general strengthening or a condition-specific sharpening of the motor representations. We used the TMS methodology to probe how context modulated activation in the motor cortex during the reading of verbs. Participants were instructed to read manualaction or non-manual verbs that appeared either in a rich sentential context (full sentence) or in a minimal context (pronoun-verb pair). TMS pulses were delivered over the pointer finger area of the left primary motor cortex at three latencies after verb presentation (200, 300, or 400 ms) to better capture individual corticospinal modulation during reading. We expect to replicate the effect of action often seen in the literature, characterized by a greater corticospinal excitability increase for action compared to non-action sentences. However, this effect may be modulated by sentence context. The strengthening idea would predict greater corticospinal excitability in the rich sentential context than in the minimal context for both manual and non-manual verbs. The sharpening idea, on the other hand, would predict an intensified action effect in the rich context, with an increase of corticospinal excitability for manual verbs but not for non-manual verbs.

Material and method

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

Participants

Thirty-five healthy right-handed adults (12 women; mean age = 23.79 years-old; range 18-29 years) participated in the experiment at the laboratory. As context may have only a subtle effect on the motor system, this sample size was chosen based on a power analysis using a small effect size of 0.25 (G*Power, version 3.1.9.2., Faul et al., 2007). Handedness was assessed by the Edinburgh inventory (Oldfield, 1971). All participants had normal or corrected-to-normal vision, without neurological, physical and cognitive pathologies. The experiment was conducted in two sessions:

1) an experimental TMS recording and 2) a reading comprehension assessment. Volunteers were first approved for participation by a medical doctor, and confirmed their participation with written consent. The Ethics Committee approved experimental protocol (CPP SOOM III, ClinicalTrials.gov Identifier: NCT03334526) and procedures in accordance with the Declaration of Helsinki.

Stimuli

One hundred and forty-four French verbs were generated; half referred to Manual Action (e.g., "I scrub") and half were Non-Manual (including non-action as "I ignore" as well as actions that did not implicate the hand as "I kick"). These verbs were presented either in a Rich context that helped to specify the verb (e.g., "I see a stain on my shirt and I scrub it" or "I see a stain on my shirt and I ignore it"), or in Minimal context (e.g., "I scrub" or "I ignore"; see Table 1 for details). Sentences always appeared in the first-person present tense. All rich-context sentences were created such that the target verb occurred at the end of the sentence (see Figure 1). This final pronoun-verb segment was presented alone on a subsequent screen after the beginning of the sentence was presented, thus yielding an identical or very similar presentation screen to the minimal sentence version.

A list was generated, in which 36 of the manual action verbs, and 36 of the non-manual verbs occurred within rich sentence contexts, while the other 36 manual action verbs, and 36 non-manual verbs occurred in minimal sentence context. A second list presented these same verbs in the opposite rich and minimal context versions. Each manual action verb (to scrub) was paired with a non-manual verb (to ignore), sharing the same rich-context sentence (I see a stain on my shirt and I scrub/ignore it), and each verbal stimulus only appeared once in the list. Thereby, across participants, each verb appeared in both conditions, but a given participant saw each verb and each sentence only once. For example, for the scrub/ignore pair, if a given participant read the minimal sentence "I scrub", she/he would also read the rich context sentence "I see a stain on my shirt and I ignore it". And another participant would read "I ignore" as well as "I see a stain on my shirt and I scrub it". In addition, TMS latencies were counterbalanced within each list.

Latent Semantic Analysis (LSA: Isa.colorado.edu; Landauer et al., 1998) was used to assure that manual action verbs (0.294±0.257) and non-manual verbs (0.295±0.269) did not significantly differ in their association scores to the preceding sentence fragments (P=0.868). We also compared various psycholinguistic factors (written frequency, number of characters, and spelling neighbors) between manual action verbs and non-manual verbs using the Lexique.org database (New et al., 2004). Corrected t-tests (modified Hochberg's procedure; Rom, 1990) yielded a significant difference only for written frequency (p=0.004; Mean ratios: 11.34±16.69 and 30.80±53.70 for manual action verbs and non-manual verbs, respectively), but not for number of characters (p=0.604; Mean ratios: 6.37±1.63 and 6.51±1.57 for manual action verbs and non-manual verbs, respectively) nor spelling neighbors (p=0.136; Mean ratios: 5.95±5.03 and 4.73±4.74 for manual action verbs and non-

manual verbs, respectively). The frequency difference is largely due to a few high-frequency non-manual verbs such as "to read." Furthermore, as higher frequency would typically increase facilitation and excitability, this difference could only hurt the chances of observing the predicted effect of action. To confirmed the accuracy of the manual and non-manual distinctions of the stimuli, 8 participants were asked to rate on a scale from 1-5 whether a sentence described a manual action or not (1= absolutely not a hand action, and 5= entirely a hand action). A t-test revealed a significant difference (p<0.001) between the manual (4.36±0.34) and non-manual (1.43±0.54) verbs.

Table 1. Example of lexical stimuli used in the experiment. In grey lines are the English translation of the French verbs/sentences presented.

Man	Manual action		Non-manual	
Minimal Context	Rich Context	Minimal	Rich context	
		context		
Je mélange	Les cartes sont sur la	Je memorise	Les cartes sont sur la	
	table et je les mélange		table et je les mémorise	
I shuffle	The cards are on the	I memorize	The cards are on the	
	table and I shuffle them		table and I memorize them	
Je frotte	Je vois une tache sur	J'ignore	Je vois une tache sur	
	mon t-shirt et je la frotte		mon t-shirt et je l'ignore	
l scrub	I see a stain on my	I ignore	I see a stain on my shirt	
	shirt and I scrub it		and I ignore it	
Je ferme	L'enveloppe est prête	Je relis	L'enveloppe est prête et	
	et je la ferme		je la relis	
l close	The envelope is ready	I reread	The envelope is ready	
	and I am closing it		and I reread it	
J'accorde	La nouvelle guitare est	J'écoute	La nouvelle guitare est	
	très belle et je l'accorde		très belle et je l'écoute	
I tune	The new guitar is very	l listen	The new guitar is very	
	beautiful and I tune it		beautiful and I listen to it	

Je paraphe	Après avoir lu le	Je contemple	Après avoir lu le contrat,
	contrat, je le paraphe		je le contemple
l initial	After reading the	I contemplate	After reading the
	contract, I initial it		contract, I contemplate it

Procedure

Participants sat in an armchair. Stimuli were presented on a 19-inch LCD monitor by a home-made software, which controlled TMS triggering and synchronized physiological recordings. Throughout the recording, participants were instructed not to move, while they silently read sentences (passive reading). Corticospinal excitability was recorded at rest, as well as during the reading task at various stimulation latencies (200, 300 or 400ms after the verb onset). These latencies were chosen based on electroencephalogram studies showing 200ms as a minimal latency for semantic processing in anterior regions (Pulvermüller et al., 2001; Hauk et al., 2006). In addition, later latencies are related to changes of P300 in parietal and frontal regions (Pulvermüller et al., 2001), but also N400 in posterior regions (Kellenbach et al., 2002; Beres, 2017) when participants read action words compared to abstract words. In the present study, these different latencies were used to account for individual differences in the speed of semantic integration, which would yield variable peaks of motor cortex involvement.

Before the experimental trials, the participants performed four familiarization trials, each with a fixation cross, followed by a minimal sentence or a rich sentential context, then a black screen before the next trial. Once the participant understood this procedure, four new practice trials were presented, now including TMS pulses after the appearance of the target verb. Then, the experimental session was divided

into 3 blocks, each comprised of 56 trials, yielding 168 trials total in the experiment. Among the 56 trials in each block, 8 trials were interspersed with TMS pulses only at rest (fixation cross), which served as reference stimulations and allowed comparisons across experimental conditions in that block. Therefore, participants saw 48 experimental trials in each of the three blocks, and 144 trials in total. Moreover, to avoid mental fatigue and lack of concentration, there was a two-minute break between each block, allowing participants to be as attentive as possible during experimental blocks. In each block, the number of stimulations was equally distributed between all conditions, but also between the three different stimulation latencies.






Figure 1.Experimental procedure. Sentences in a Minimal or Rich context were presented on a screen in front of the participant. Single-pulse TMS was triggered over the left hemisphere and motor-evoked potentials recorded in the right index finger.

Several weeks later, individuals participated in a second experimental session, in which two levels of contextual reading comprehension ability were assessed. This session was included in order to account for the possible effect of individual differences in the use of context during reading comprehension. To investigate sensitivity to lower-level memory-based context, we administered a "Remember versus Know" paradigm (Tulving, 1985), in which they had to identify whether words were encountered in a previously read text (yes or no response), and in what context they were encountered. To assess higher-level use of context during comprehension, we used a Curriculum-Based Measurement-Maze (Parker et al., 1992), in which readers rely on inferences to select a contextually-appropriate word from three alternatives. After the first sentence, every seventh word in the passage is replaced with that correct word and two distractors. Participants must use the context to select the word that fits best with the rest of the passage.

TMS

Single-pulse TMS was generated from an electromagnetic stimulator Magstim 200 (Magstim Company Ltd, Whitland) and using a figure-of-eight coil (70 mm in diameter). The coil was placed over the contralateral left hemisphere to target the motor area of the First Dorsal Interosseous (FDI) muscle of the right hand. The coil rested tangential to the scalp with the handle pointing backwards and laterally at a 45° angle away from the midline. First, the mapping of the motor cortex of each participant determined the precise stimulation site (hotspot) corresponding to the location, where the MEPs amplitude of the FDI muscle was the highest and the most consistent for the same stimulation intensity. The resting motor threshold of each participant was determined as the minimal intensity of TMS necessary to induce a MEP of 50µV peak-to-peak amplitude in the right FDI muscle for 4 trials out of 8.

During the experimental session, TMS intensity was set at 130% of the resting motor threshold. This intensity was chosen not only to stimulate the participant above the motor threshold, but also to achieve a stimulus intensity near the middle of the stimulus-response curve (Groppa et al., 2012), from which point it would be possible to observe both increases and decreases in MEP amplitude

EMG recording

The EMG signal was recorded through 10mm-diameter surface electrodes (Contrôle Graphique Médical, Brice Comte-Robert, France) placed over the FDI muscle of the right hand. Before placing the electrodes, the skin was shaved and cleaned in order to reduce noise in the EMG signal (< 20µV). The EMG signals were amplified and bandpass filtered on-line (10-1000 Hz, Biopac Systems Inc.) and digitized at 2000 Hz for off-line analysis. We also measured the root mean square of EMG signal (EMGrms), to ensure that participants were at rest in all conditions. Throughout the experiment, EMGrms was monitored for the 100 ms preceding every TMS pulse to ensure complete muscle relaxation, using the following formula (MD = movement duration):

$$RMS = \sqrt{rac{1}{MD}\int\limits_{0}^{MD}{(EMG)^{2}dt}}$$

Data and statistical analysis

EMG data were extracted with Matlab (The MathWorks, Natick, Massachusetts, USA) and we measured peak-to-peak MEP amplitude. We first eliminated outliers for each participant within each stimulation latency and condition; data falling 2 SDs

above or below individual means for each experimental condition were removed before analysis (4.13% of total). Then, the average MEP amplitude for each condition was normalized with reference to baseline MEP amplitude (rest condition):

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

$$\left(\frac{Mean\ condition-Mean\ rest}{Mean\ rest}\right) X\ 100$$

Importantly, no analysis on the timing factor was performed as we rather isolated the highest average amplitude (over 12 trials) among the three stimulation times (200, 300, or 400ms) for each participant in each condition. Thus, for each participant, we use the data from only one of these three latencies for each condition, presumably corresponding to the moment when the motor representation was most active. Individuals can vary greatly in speed of decoding, as well as lexical access (Sereno et al., 1998; Dambacher et al., 2006) and semantic integration (Kutas and Hillyard, 1980; Brown and Hagoort, 1993; Van Berkum et al., 1999). In addition, the motor activation that we aimed to measure is likely punctual and fleeting in nature. Therefore, this method of stimulation at multiple latencies after the target verb (informed by previous investigations) was designed to maximize the chances of tapping into the fleeting motor representation, which may might not even be present at all latencies. Selecting only the latency with the highest average MEP helps to account for inter-individual variability in the latency of the motor representation, and its consequential increase in corticospinal excitability. Finally, Grubbs test identified extreme mean values for one participant, who was excluded from the final analysis. All data analyses were performed using the software Statistica (Stat Soft, France).

Data normality was confirmed by the Shapiro-Wilk test. To assess the influence of *Action* and *Context* on corticospinal excitability, we performed a 2 by 2 repeated measures ANOVA with *Action* (Manual action vs. Non-manual) and *Context* (Minimal vs. Rich) as within-subject factors. Then, one-sample t-tests were used to compare

normalized MEPs at zero for each condition to assess whether corticospinal excitability during reading changed from baseline. Finally, in order to ensure that our results were not contaminated by muscular pre-activity, we tested with Wilcoxon tests if the EMGrms before the TMS artifact was not different from zero in our experimental conditions.

To assess the influence of participants' comprehension ability on their corticospinal excitability modulations by context, we performed Pearson correlations between the context effect (Minimal vs. Rich) and the memory-based, as well as the higher context-level comprehension scores. The data are presented as mean values (±standard error) and the alpha value was set at 0.05.

3. Results

 The overall ANOVA revealed a main effect of *Action* ($F_{1,33}$ =4.811, p=0.035, ηp^2 =0.127), with larger MEP ratios for manual action verbs (11.02 ±3.25%) than non-manual verbs (6.76 ±3.19%). We did not observe a main effect of *Context* ($F_{1,33}$ =0.168, p=0.683, ηp^2 =0.005) nor an *Action* by *Context* interaction ($F_{1,33}$ =0.064 p=0.801, ηp^2 =0.001) (Figure 2).





Figure 2.Corticospinal excitability in Rich and Minimal Context. Box plots on the left side represent normalized MEPs. The line represents the mean. The box and the whiskers represent the 25th/75th quartile and the 5th and 95th

Using one-sample t-tests, we found that normalized MEPs of all conditions were different from baseline (all p's<0.05), except for non-manual verbs in the rich context (p=0.167). This finding reveals the possible importance of context for reducing residual activation for non-manual verbs that may be unclear in the absence of sentence context.

Low and high-level reading comprehension abilities influenced these outcomes, and in particular the context effect. We observed a positive correlation between the modulations of corticospinal excitability in rich context conditions and the low-level reading comprehension scores (p=0.048; r=0.341). Specifically, higher scores in memory-based comprehension yielded greater corticospinal excitability in rich context conditions but not minimal (p=0.495; r=0.121). This finding reveals that the modulation of corticospinal excitability varies across individuals, and that comprehension ability may partially explain such variability. No significant correlations were observed for the high-level context-based comprehension scores (minimal context: p=0.623; r=-0.087; rich context: p=0.590; r=0.095) (Figure 3). The four correlations were evaluated using the modified Hochberg's correction procedure (Rom, 1990).

Memory-based comprehension scores



Higher context-based comprehension scores




Figure 3. Correlations between the corticospinal excitability modulations depend on the context (Minimal vs. Rich) and the comprehension scores (memory vs. context-level). (A and B) represent correlations between the excitability modulations in Minimal context (A) and Rich context (B) with the memory-based comprehension scores. (C and D) represent correlations between the excitability modulations in Minimal context (C) and Rich context (D) with the higher context-level comprehension scores.

Finally, we analyzed the EMGrms prior to TMS artifacts to test for confounds in MEP modulation. We did not observe any difference between baseline (mean \pm SD = 1.76 \pm 1.06 μ V) and the other conditions (all p's>0.05; in minimal context, for manual verbs: 1.83 \pm 1.10 μ V, for non-manual verbs: 1.84 \pm 1.13 μ V; in rich context, for manual verbs: 1.90 \pm 1.48 μ V, for non-manual verbs: 1.83 \pm 1.11 μ V). This indicates that MEP modulations were not influenced by background EMGrms.

Discussion

The results of this study support the importance of the type of stimuli in modulating the involvement of the motor cortex during action language processing. We observed action-specific modulation of corticospinal excitability, with increased MEP amplitude for manual action verbs compared to non-manual verbs. Nonetheless, we did not observe a main effect of context nor an interaction between action and context. These findings do not support the "strengthening" idea, whereby sentence context would generate greater motor activation in general. However, there does seem to be some evidence for the idea of "sharpening," as motor activation did not differ from rest when the verb phrase was presented in combination with a sentence context that makes it clear that there is no manual action.

The activation of the motor cortex is presumed to arise from reading events describing actions, typically verbs. However, even a non-action verb may be able to evoke some degree of motor activation if it is underspecified, because the event representation is vague. Indeed, it seems odd that non-manual verbs presented in a minimal context might increase the corticospinal excitability in hand muscles. However, in the absence of context, it is possible that lexical access to the less constrained meaning evokes a broad distributed activation, including the motor system for some words. Additionally, our non-manual-action verbs included actions that were not manual, and when presented out of context, perhaps these could broadly activate the motor system, including the hand area. Contextual constraints may reduce the possibilities for such broad activation and eliminate unintended meanings. If the verb is well contextualized, the motor representation is more precise, yielding activation of only the appropriate effectors in the motor system that are relevant to the event. This is reflected in our results where all conditions resulted in a higher state of corticospinal excitability from the baseline, except for non-manual

verbs in the rich context. This suggests that the motor cortex can be automatically activated at some level even by non-action verbs in the absence of context [see also motor activation in abstract verbs (Sakreida et al., 2013) and metaphoric sentences (Desai et al., 2013; Reilly et al., 2019)].

While we did not observe any boost in motor activation when a rich context specifies the nature of the described situation, we did observe a lack of residual motor activation from non-manual verbs for which the representation may have been unclear in the absence of context. The rich context may lead to a disambiguation of the meaning of the verb, especially when the verb is vague or has multiple meanings. This would not apply to all of our stimuli, as for some verb phrases, it was already clear whether a manual action was implicated or not, even without the context. However, at least some of our non-manual stimuli were interpreted as implicating less manual action when presented in combination with a receding sentence context. Consistent with Mahon and colleagues (Mahon and Caramazza, 2008; Mahon, 2015), motor cortex activation may therefore play an important role in embellishing certain aspects of word meaning that are relevant in a given context.

Moreover, an action effect in minimal context is consistent with previous studies (Papeo et al., 2009; Innocenti et al., 2014), reporting that many manual action verbs would activate the motor cortex to a greater extent than non-manual verbs without context. However, we expected a greater difference for these same stimuli if accompanied by more contextual information to better specify the action, supporting the sharpening idea. Indeed, the manual nature of the meaning of some verbs is quite precise regardless of context (e.g., "I write") whereas others can be rather vague (e.g., "I take"). As mentioned above, this vagueness can even introduce motor cortex activation in verbs that are intended to be non-action stimuli. Although we did

not observe a greater effect of action in the rich compared minimal context, further research is currently underway to further test this idea of sharpening motor activation.

The present study suggests that the context effect is sensitive to individual factors, such as memory-based comprehension ability. We found that participants with higher comprehension ability exhibited a greater increase in excitability when reading rich-context sentences, regardless of verb type. This does not necessarily mean that context automatically strengthens motor activity (since no main effect of context was observed), but rather that the engagement of the motor cortex can vary across situations and across individuals, and may depend on parameters such as comprehension ability.

Based on the current findings and previous studies (Zwaan et al., 2010; Van Dam et al., 2012; van Dam et al., 2012; Gilead et al., 2013; Van Dam et al., 2014; Beauprez et al., 2018, 2020; Wurm and Caramazza, 2019), language-induced motor representations are flexible and at least to some extent sensitive to linguistic context. It appears that context may play a role in strengthening motor activation for good comprehenders, as well as eliminating residual motor activation for non-action verbs. These findings are consistent with cognitive theories that accommodate semantic flexibility and top-down processes, whereby relevant features and alternative meanings are determined by the semantic context very early as each new word is encountered (e.g., Gentner, 1981; Marslen-Wilson, 1987; Tomasino and Rumiati, 2013).

The question remains as to whether activation of the primary motor cortex is necessary to process action language (see also: Wurm and Caramazza, 2019). Researchers observed that disturbances of motor areas caused by repeated TMS

during language processing can yield a decrease in performance, implying the engagement of the motor cortex in this process (Willems et al., 2011; Repetto et al., 2013; Vukovic et al., 2017; Courson et al., 2018). While understanding might be disturbed, diminished, or less effective, it is not completely blocked. Although language comprehension may be able to proceed without the involvement of the primary motor cortex, our findings suggest that it may play a role in the optimization of this process, refining linguistic understanding and making it more effective when necessary.

One potential limitation of the current study is the number of experimental trials per participant. Since we cannot know exactly when the motor representation will engage and reach its maximum for each subject, we have to probe at several latencies on separate trials, and select the peak afterwards. This can be considered a strength of our study in that we consider individual differences in processing time, but it can also be a limitation in that we reduce the number of experimental trials. In future investigations, we plan to overcome this limitation by calibrating each subject's individual processing time prior to the test session, as well as employing continuous measures, such as EEG. Finally, while we strove to maximize the control between the rich and minimal context conditions, we could not avoid the additional pronoun which sometimes preceded the verb in the rich context condition. When this occurred, the target verb was rendered grammatically transitive in the rich context (*Je frotte*; I scrub it) while it remained intransitive in the minimal context (*Je frotte*; I scrub). Otherwise, the probe screens were identical for the two conditions.

In conclusion, this study provides relevant information about the role of sentential context on primary motor cortex activation. We found that linguistic context serves to focus the motor representation of the described action, eliminating residual

motor activation for non-action verbs. Moreover, the context effect appears to be sensitive to individual differences in reading comprehension. Consistent with the idea of semantic flexibility and top-down processes, when target verb phrases are presented with a sentence context that makes it clear that no manual action is evoked, the motor cortex does not contribute to the meaning representation.

Declaration of competing interest

None declared.

457

458

459

460

461

462

464

Author Contributions

- 465 Experiment design: WD, CML, FL, CP
- 466 Data collection: WD
- 467 Statistical analysis: WD, CML, FL
- 468 Manuscript preparation: WD, CP, CML, FL

Data availability statement.

- 470 All data from this study are available at
- 471 https://osf.io/5zbmq/?view_only=82c93af470fd42be8ca33969b14f1c37

472 Acknowledgement

- The authors are very grateful to Cyril Sirandré for help with the custom

 Neurostim software and the experimental setup.
- 475 **References**
- Andres M, Finocchiaro C, Buiatti M, Piazza M (2015) Contribution of motor representations to action verb processing. Cognition 134:174–184.

- Aziz-Zadeh L, Wilson SM, Rizzolatti G, Iacoboni M (2006) Congruent Embodied
 Representations for Visually Presented Actions and Linguistic Phrases Describing
- 480 Actions. Current Biology 16:1818–1823.
- Barsalou LW (1999) Perceptual symbol systems. Behavioral and Brain
- 482 Sciences 22:577–609.
- Barsalou LW (2008) Grounded Cognition. Annual Review of Psychology
- 484 59:617–645.
- Barsalou LW, Simmons WK, Barbey AK, Wilson CD (2003) Grounding
- 486 conceptual knowledge in modality-specific systems. Trends in Cognitive Sciences
- 487 **7:84–91**.
- Beauprez SA, Blandin Y, Almecija Y, Bidet-Ildei C (2020) Physical and
- 489 observational practices of unusual actions prime action verb processing. Brain and
- 490 Cognition 138.
- Beauprez SA, Toussaint L, Bidet-Ildei C (2018) When context modulates the
- influence of action observation on language processing. PLoS ONE 13.
- Beres AM (2017) Time is of the Essence: A Review of Electroencephalography
- 494 (EEG) and Event-Related Brain Potentials (ERPs) in Language Research. Applied
- 495 Psychophysiology Biofeedback 42:247–255.
- Brown C, Hagoort P (1993) The processing nature of the N400: Evidence from
- 497 masked priming. Journal of Cognitive Neuroscience 5:34–44.
- 498 Buccino G, Riggio L, Melli G, Binkofski F, Gallese V, Rizzolatti G (2005)
- 499 Listening to action-related sentences modulates the activity of the motor system: A
- combined TMS and behavioral study. Cognitive Brain Research 24:355–363.
- Courson M, Macoir J, Tremblay P (2018) A facilitating role for the primary motor
- 502 cortex in action sentence processing. Behavioural Brain Research 336:244–249.
- 503 Dalla Volta R, Gianelli C, Campione GC, Gentilucci M (2009) Action word
- understanding and overt motor behavior. Experimental Brain Research 196:403–412.

- Dambacher M, Kliegl R, Hofmann M, Jacobs AM (2006) Frequency and predictability effects on event-related potentials during reading. Brain Research 1084:89–103.
- de Zubicaray G, Arciuli J, McMahon K (2013) Putting an "end" to the motor cortex representations of action words. Journal of cognitive neuroscience 25:1957–1974.
- Decety J, Grèzes J (2006) The power of simulation: Imagining one's own and other's behavior. Brain Research 1079:4–14.
- Desai RH, Conant LL, Binder JR, Park H, Seidenberg MS (2013) A piece of the action: Modulation of sensory-motor regions by action idioms and metaphors.

 Neurolmage 83:862–869.
- Faul F, Erdfelder E, Lang A-G, Buchner A (2007) G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. Behav Res Methods 39:175–191.
- Fischer MH, Zwaan RA (2008) Embodied language: A review of the role of the motor system in language comprehension. Quarterly Journal of Experimental Psychology 61:825–850.
- Franklin ZC, Wright DJ, Holmes PS (2020) Using action-congruent language facilitates the motor response during action observation: A combined transcranial magnetic stimulation and eye-tracking study. Journal of Cognitive Neuroscience 32:634–645.
- Gallese V, Lakoff G (2005) The brain's concepts: The role of the sensory-motor system in conceptual knowledge. Cognitive Neuropsychology 22:455–479.
- Gentilucci M, Benuzzi F, Bertolani L, Daprati E, Gangitano M (2000) Language and motor control. Experimental Brain Research 133:468–490.
- Gentner D (1981) Verb semantic structures in memory for sentences: Evidence for componential representation. Cognitive Psychology 13:56–83.
- Gianelli C, Kühne K, Lo Presti S, Mencaraglia S, Dalla Volta R (2020) Action processing in the motor system: Transcranial Magnetic Stimulation (TMS) evidence

- of shared mechanisms in the visual and linguistic modalities. Brain and Cognition 139.
- Gilead M, Liberman N, Maril A (2013) The language of future-thought: An fMRI study of embodiment and tense processing. NeuroImage 65:267–279.
- Glenberg AM, Gallese V (2012) Action-based language: A theory of language acquisition, comprehension, and production. Cortex 48:905–922.
- Groppa S, Oliviero A, Eisen A, Quartarone A, Cohen LG, Mall V, Kaelin-Lang A, Mima T, Rossi S, Thickbroom GW, Rossini PM, Ziemann U, Valls-Solé J, Siebner HR (2012) A practical guide to diagnostic transcranial magnetic stimulation: Report of an IFCN committee. Clinical Neurophysiology 123:858–882.
- Grossman M, Koenig P, DeVita C, Glosser G, Alsop D, Detre J, Gee J (2002)
 Neural representation of verb meaning: An fMRI study. Human Brain Mapping
 15:124–134.
- Hauk O, Davis MH, Ford M, Pulvermüller F, Marslen-Wilson WD (2006) The time course of visual word recognition as revealed by linear regression analysis of ERP data. Neurolmage 30:1383–1400.
- Hauk O, Johnsrude I, Pulvermüller F (2004) Somatotopic Representation of Action Words in Human Motor and Premotor Cortex. Neuron 41:301–307.
- Innocenti A, De Stefani E, Sestito M, Gentilucci M (2014) Understanding of action-related and abstract verbs in comparison: a behavioral and TMS study. Cognitive processing 15:85–92.
- Jeannerod M (2008) Motor Cognition: What Actions Tell the Self.
- Kellenbach ML, Wijers AA, Hovius M, Mulder J, Mulder G (2002) Neural differentiation of lexico-syntactic categories or semantic features? Event-related potential evidence for both. Journal of Cognitive Neuroscience 14:561–577.
- Kemmerer D (2015) Are the motor features of verb meanings represented in the precentral motor cortices? Yes, but within the context of a flexible, multilevel architecture for conceptual knowledge. Psychonomic Bulletin and Review 22:1068–1075.

- Klepp A, van Dijk H, Niccolai V, Schnitzler A, Biermann-Ruben K (2019) Action verb processing specifically modulates motor behaviour and sensorimotor neuronal oscillations. Scientific reports 9:15985.
- Kutas M, Hillyard SA (1980) Reading senseless sentences: Brain potentials reflect semantic incongruity. Science 207:203–205.
- Labruna L, Fernández-Del-Olmo M, Landau A, Duqué J, Ivry RB (2011)
- Modulation of the motor system during visual and auditory language processing.
- 570 Experimental Brain Research 211:243–250.
- Lambon Ralph MA, Pobric G, Jefferies E (2009) Conceptual knowledge is
- 572 underpinned by the temporal pole bilaterally: Convergent evidence from rTMS.
- 573 Cerebral Cortex 19:832–838.
- Lambon Ralph MA, Sage K, Jones RW, Mayberry EJ (2010) Coherent concepts
- are computed in the anterior temporal lobes. Proceedings of the National Academy of
- 576 Sciences of the United States of America 107:2717–2722.
- Landauer TK, Foltz PW, Laham D (1998) An introduction to latent semantic
- analysis. Discourse processes 25:259–284.
- Longe O, Randall B, Stamatakis EA, Tyler LK (2007) Grammatical categories in
- the brain: The role of morphological structure. Cerebral Cortex 17:1812–1820.
- 581 Mahon BZ (2015) What is embodied about cognition? Language, Cognition and
- 582 Neuroscience 30:420–429.
- 583 Mahon BZ, Caramazza A (2008) A critical look at the embodied cognition
- 584 hypothesis and a new proposal for grounding conceptual content. Journal of
- 585 Physiology Paris 102:59–70.
- Marslen-Wilson WD (1987) Functional parallelism in spoken word-recognition.
- 587 Cognition 25:71–102.
- New B, Pallier C, Brysbaert M, Ferrand L (2004) Lexique 2: A new French
- lexical database. Behavior Research Methods, Instruments, and Computers 36:516-
- 590 524.

- Oldfield RC (1971) The assessment and analysis of handedness: The Edinburgh inventory. Neuropsychologia 9:97–113.
- Papeo L, Lingnau A, Agosta S, Pascual-Leone A, Battelli L, Caramazza A (2015) The origin of word-related motor activity. Cerebral Cortex 25:1668–1675.
- Papeo L, Pascual-Leone A, Caramazza A (2013) Disrupting the brain to validate hypotheses on the neurobiology of language. Frontiers in Human Neuroscience 7.
- Papeo L, Rumiati RI, Cecchetto C, Tomasino B (2012) On-line changing of thinking about words: The effect of cognitive context on neural responses to verb reading. Journal of Cognitive Neuroscience 24:2348–2362.
- Papeo L, Vallesi A, Isaja A, Rumiati RI (2009) Effects of TMS on different stages of motor and non-motor verb processing in the primary motor cortex. PloS one 4:e4508.
- Parker R, Hasbrouck JE, Tindal G (1992) The Maze as a Classroom-Based Reading Measure: Construction Methods, Reliability, and Validity. The Journal of Special Education 26:195–218.
- Patterson K, Nestor PJ, Rogers TT (2007) Where do you know what you know?

 The representation of semantic knowledge in the human brain. Nature Reviews

 Neuroscience 8:976–987.
- Perani D, Cappa SF, Schnur T, Tettamanti M, Collina S, Rosa MM, Fazio F (1999) The neural correlates of verb and noun processing a PET study. Brain 122:2337–2344.
- Pobric G, Jefferies E, Lambon Ralph MA (2010) Amodal semantic representations depend on both anterior temporal lobes: Evidence from repetitive transcranial magnetic stimulation. Neuropsychologia 48:1336–1342.
- Pulvermüller F, Härle M, Hummel F (2001) Walking or talking?: Behavioral and neurophysiological correlates of action verb processing. Brain and Language 78:143–168.

- Pulvermüller F, Hauk O, Nikulin V V., Ilmoniemi RJ (2005) Functional links between motor and language systems. European Journal of Neuroscience 21:793– 797.
- Rabahi T, Fargier P, Rifai Sarraj A, Clouzeau C, Massarelli R (2013) Effect of Action Verbs on the Performance of a Complex Movement. PLoS ONE 8:e68687.
- Rabahi T, Sarraj AR, Fargier P, Clouzeau C, Massarelli R (2012) Action verb and motor performance. Kinesitherapie 12:42–46.
- Raposo A, Moss HE, Stamatakis EA, Tyler LK (2009) Modulation of motor and premotor cortices by actions, action words and action sentences. Neuropsychologia 47:388–396.
- Reilly M, Howerton O, Desai RH (2019) Time-course of motor involvement in literal and metaphoric action sentence processing: A TMS study. Frontiers in Psychology 10.
- Repetto C, Colombo B, Cipresso P, Riva G (2013) The effects of rTMS over the primary motor cortex: The link between action and language. Neuropsychologia 51:8–13.
- Rogers TT, Hocking J, Noppeney U, Mechelli A, Gorno-Tempini ML, Patterson K, Price CJ (2006) Anterior temporal cortex and semantic memory: Reconciling findings from neuropsychology and functional imaging. Cognitive, Affective and Behavioral Neuroscience 6:201–213.
- Rom DM (1990) A sequentially rejective test procedure based on a modified bonferroni inequality. Biometrika 77:663–665.
- Rüschemeyer SA, Brass M, Friederici AD (2007) Comprehending prehending:
 Neural correlates of processing verbs with motor stems. Journal of Cognitive
 Neuroscience 19:855–865.
- Sakreida K, Scorolli C, Menz MM, Heim S, Borghi AM, Binkofski F (2013) Are abstract action words embodied? An fMRI investigation at the interface between language and motor cognition. Frontiers in human neuroscience 7:125.

- Scorolli C, Jacquet PO, Binkofski F, Nicoletti R, Tessari A, Borghi AM (2012)

 Abstract and concrete phrases processing differentially modulates cortico-spinal

 excitability. Brain Research 1488:60–71.
- Sereno SC, Rayner K, Posner MI (1998) Establishing a time-line of word recognition: Evidence from eye movements and event-related potentials.

 NeuroReport 9:2195–2200.
- Taylor LJ, Zwaan RA (2008) Motor resonance and linguistic focus. Quarterly
 Journal of Experimental Psychology 61:896–904.
- Tomasino B, Rumiati RI (2013) At the Mercy of Strategies: The Role of Motor Representations in Language Understanding. Frontiers in Psychology 4.
- Tulving E (1985) Memory and consciousness. Canadian Psychology/Psychologie canadienne 26:1–12.
- Van Berkum JJA, Hagoort P, Brown CM (1999) Semantic integration in sentences and discourse: Evidence from the N400. Journal of Cognitive Neuroscience 11:657–671.
- Van Dam WO, Brazil IA, Bekkering H, Rueschemeyer SA (2014) Flexibility in embodied language processing: Context effects in lexical access. Topics in Cognitive Science 6:407–424.
- Van Dam WO, Desai RH (2017) Embodied Simulations Are Modulated by Sentential Perspective. Cognitive science 41:1613–1628.
- Van Dam WO, Rueschemeyer SA, Bekkering H (2010) How specifically are action verbs represented in the neural motor system: An fMRI study. NeuroImage 53:1318–1325.
- van Dam WO, Rueschemeyer SA, Lindemann O, Bekkering H (2010) Context effects in embodied lexical-semantic processing. Frontiers in Psychology 1.
- Van Dam WO, Van Dijk M, Bekkering H, Rueschemeyer SA (2012) Flexibility in embodied lexical-semantic representations. Human Brain Mapping 33:2322–2333.

- van Dam WO, van Dongen E V., Bekkering H, Rueschemeyer SA (2012)
 Context-dependent changes in functional connectivity of auditory cortices during the
 perception of object words. Journal of Cognitive Neuroscience 24:2108–2119.
- Vukovic N, Feurra M, Shpektor A, Myachykov A, Shtyrov Y (2017) Primary motor cortex functionally contributes to language comprehension: An online rTMS study. Neuropsychologia 96:222–229.
- Willems RM, Labruna L, D'Esposito M, Ivry R, Casasanto D (2011) A Functional Role for The Motor System in Language Understanding: Evidence from Theta-Burst Transcranial Magnetic Stimulation. Psychological Science 22:849–854.
- 683 Wurm MF, Caramazza A (2019) Distinct roles of temporal and frontoparietal 684 cortex in representing actions across vision and language. Nature Communications 685 10.
- Xu X, Kang C, Guo T (2016) Imageability and semantic association in the representation and processing of event verbs. Cognitive Processing 17:175–184.
- Zwaan RA, Taylor LJ (2006) Seeing, acting, understanding: Motor resonance in language comprehension. Journal of Experimental Psychology: General 135:1–11.
- Zwaan RA, Taylor LJ, de Boer M (2010) Motor resonance as a function of narrative time: Further tests of the linguistic focus hypothesis. Brain and Language 112:143–149.
- Zwaan RA, van der Stoep N, Guadalupe T, Bouwmeester S (2012) Language comprehension in the balance: the robustness of the action-compatibility effect (ACE). PloS one 7:e31204.

696