

Influence of Voluntary Contraction Level, Test Stimulus Intensity and Normalization Procedures on the Evaluation of Short-Interval Intracortical Inhibition

Cécilia Neige, Sidney Grosprêtre, Alain Martin, Florent Lebon

► To cite this version:

Cécilia Neige, Sidney Grosprêtre, Alain Martin, Florent Lebon. Influence of Voluntary Contraction Level, Test Stimulus Intensity and Normalization Procedures on the Evaluation of Short-Interval Intracortical Inhibition. Brain Sciences, 2020, 10 (7), 10.3390/brainsci10070433. hal-04185348

HAL Id: hal-04185348 https://u-bourgogne.hal.science/hal-04185348

Submitted on 22 Aug 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Article



Influence of Voluntary Contraction Level, Test Stimulus Intensity and Normalization Procedures on the Evaluation of Short-Interval Intracortical Inhibition

Cécilia Neige 1,*, Sidney Grosprêtre ², Alain Martin ¹ and Florent Lebon ¹

- ¹ INSERM UMR1093-CAPS, Université Bourgogne Franche-Comté, UFR des Sciences du Sport, F-21078 Dijon, France; alain.martin@u-bourgogne.fr (A.M.); florent.lebon@u-bourgogne.fr (F.L.)
- ² EA4660-C3S Laboratory Culture, Sport, Health and Society, Université Bourgogne Franche-Comté,
- 25000 Besançon, France; sidney.grospretre@univ-fcomte.fr
- * Correspondence: cecilia.neige@u-bourgogne.fr; Tel.: +33-3-8039-6761

Received: 27 May 2020; Accepted: 6 July 2020; Published: 8 July 2020

Abstract: Short-interval intracortical inhibition (SICI) represents an inhibitory phenomenon acting at the cortical level. However, SICI estimation is based on the amplitude of a motor-evoked potential (MEP), which depends on the discharge of spinal motoneurones and the generation of compound muscle action potential (M-wave). In this study, we underpin the importance of taking into account the proportion of spinal motoneurones that are activated or not when investigating the SICI of the right flexor carpi radialis (normalization with maximal M-wave (Mmax) and MEPtest, respectively), in 15 healthy subjects. We probed SICI changes according to various MEPtest amplitudes that were modulated actively (four levels of muscle contraction: rest, 10%, 20% and 30% of maximal voluntary contraction (MVC)) and passively (two intensities of test transcranial magnetic stimulation (TMS): 120 and 130% of motor thresholds). When normalized to MEPtest, SICI remained unchanged by stimulation intensity and only decreased at 30% of MVC when compared with rest. However, when normalized to Mmax, we provided the first evidence of a strong individual relationship between SICI and MEPtest, which was ultimately independent from experimental conditions (muscle states and TMS intensities). Under similar experimental conditions, it is thus possible to predict SICI individually from a specific level of corticospinal excitability in healthy subjects.

Keywords: SICI; muscle contraction; transcranial magnetic stimulation; corticospinal excitability

1. Introduction

Cortical inhibitory and excitatory circuits within the primary motor cortex (M1) play an important role in the fine tuning of descending motor commands and in the neural adaptation following motor training [1]. For example, the reduction in intracortical inhibition within M1 is a crucial part of neural adaptation following acute and multi-session challenging strength training programs [2–4]. Paired-pulse transcranial magnetic stimulation (TMS) enables the non-invasive examination of these intracortical excitatory and inhibitory circuits at the time of the stimulation [5,6]. Short-interval intracortical inhibition (SICI) is a well-established paired-pulse measure to evaluate inhibitory circuits within the M1 area. SICI results from a subthreshold conditioning stimulus (CS) followed 1 to 6 ms later by a suprathreshold test stimulus (TS) applied through the same coil over M1 [5]. The high-intensity suprathreshold TS leads to a corticospinal output called motor-evoked potential (MEP), measured peripherally by means of electromyographic (EMG) recording over the target muscle. With optimal settings, the CS exerts a decrease in the conditioned MEP (MEP_{cond})

amplitude in comparison with the unconditioned MEP (MEPtest) induced by a TS alone [5]. It is thought that the CS activates low-threshold inhibitory interneurons that employ γ-aminobutyric acid type A receptor (GABAA) and reduce the excitatory inputs activated by the TS [5,7,8]. The amount of SICI is conventionally quantified by expressing the mean MEP_{cond} amplitude as a percentage of the mean MEP_{test} amplitude [5] or the difference between the mean MEP_{cond} and the mean MEP_{test} then expressed as a percentage of the mean MEP_{test} amplitude (SICI_{MEPtest}) [9]. An issue with this normalization is that the size of the MEP_{test} is known to influence SICI outcome. Indeed, previous studies have shown that in resting hand muscles, the amount of SICI_{MEPtest} increased for MEP_{test} amplitude up to 1 mV, but further remained unchanged for larger MEP_{test} amplitude, with a trend to a linear relationship between the MEP_{cond} and MEP_{test} [9–12].

Importantly, it is widely accepted that SICI represents an inhibitory phenomenon acting purely at the cortical level (rather than subcortical or spinal levels); indeed, the CS does not alter the spinal excitability when delivered at low stimulation intensities (<80% of the motor threshold) and measured in a static muscle [5,13], as evidenced by epidural recordings [7]. However, one has to keep in mind that SICI estimation is based on the MEPcond amplitude recorded peripherally, and therefore also depends critically on the discharge of spinal motoneurones and the generation of a compound muscle action potential [9,14]. Taking into account this important point, Lackmy and Marchand-Pauvert (2010) assessed SICI_{Mmax} by calculating the difference between the mean MEP_{cond} and the mean MEPtest, then expressed as a percentage of maximal compound muscle action potentials (Mmax (M-wave)). Mmax is evoked when all the motor axons of the target muscle are activated simultaneously by a supramaximal peripheral nerve stimulation. SICIMmax normalisation has the methodological benefit of considering the proportion of spinal motoneurones activated by the TS. Interestingly, Lackmy and Marchand-Pauvert (2010) found increased SICI_{Mmax} with increasing TS intensities (i.e., increased MEPtest/Mmax ratio) [9]. However, this work has only focused on resting hand muscle, whereas muscle activation is known to decrease the amount of SICIMEPtest [15,16]. To fill this gap, Opie and Semmler (2014) subsequently tested whether SICIMmax (calculated with MEPtest expressed relative to Mmax) was influenced when the muscle was voluntary activated. Contrary to SICIMEPtest, they found that the amount of SICIMmax during the active state increased concomitantly with an increase in MEPtest/Mmax ratio [14]. Importantly, larger MEPtest/Mmax ratios were obtained by increasing the TS intensity (from 110% until 150% of the motor threshold) during the active state. However, in this study, active muscle condition was assessed with only one low level of contraction (5% of maximal voluntary contraction (MVIC)), whereas SICIMEPtest has been shown to vary with increasing level of muscle contraction and MEP_{test} amplitude [10,16–19]. It is not yet known whether SICI_{Mmax} is modulated when assessed at higher levels of muscle contraction.

The aim of the present study was to broaden current knowledge on the necessity of considering the proportion of spinal motoneurones activated by single- and paired-pulse TMS in different muscle states. We quantified SICI with varying MEP_{test} amplitudes modulated actively (with four levels of muscle contraction) and passively (with two TS intensities of stimulation). The hypothesis was that SICI_{Mmax} would increase in response to larger MEP_{test}/Mmax ratios obtained by an increasing muscle contraction force. By considering this prediction as well as Opie and Semmler's (2014) findings, we also hypothesized that the amount of SICI_{Mmax} would be highly dependent upon the proportion of motoneurones activated by single-pulse TMS, regardless of the conditions leading to MEP_{test} increase.

2. Materials and Methods

2.1. Participants

Fifteen healthy volunteers (six females; mean age 24.9 years, range 22–30 years; right-handed as assessed by the Edinburgh Handedness Inventory [20]) participated in the current study after providing written informed consent. All volunteers were screened by a medical doctor for contraindications to TMS [21] and were excluded if they had a history of psychiatric or neurological illness. The protocol was approved by the "Comité de Protection des Personnes Sud-Ouest et Outre

Mer III" (CPP SOOM III) ethics committee (number 2017-A00064-49; ClinicalTrials.gov Identifier: NCT03334526) and complied with the Declaration of Helsinki.

2.2. General Experimental Procedure

Participants came to the laboratory for a single 1.5-h session and were seated in an isokinetic dynamometer chair (Biodex System 3, Biodex Medical Systems Inc., Shirley, NY, USA). The participants' right hand was firmly strapped in a neutral position to a custom-built accessory adapted for wrist movement recordings. The rotation axis of the dynamometer was aligned with the styloid process of the ulna. The upper arm was vertical along the trunk (shoulder abduction and elevation angles at 0°) and the forearm semipronated and flexed at 90° (Figure 1). Before starting the main experiment, participants familiarized themselves with the procedure of the voluntary force production feedback during an approximately 5 min warm-up of wrist flexions. They received visual feedback of all exerted force contraction on a computer screen located 1-m in front of them. Participants then performed 3 maximal voluntary isometric contractions (MVIC) lasting 3 s with at least 30 s rest in between. Verbal encouragement was provided during MVIC trials to achieve a maximal force. The maximum of the three trials was defined as the participant's MVIC. This value was then used to calculate the muscle contraction levels (i.e., 10, 20 and 30% MVIC) tested in the current study.



Figure 1. Schematic view of the experimental setup. Transcranial magnetic stimulation (TMS); median nerve stimulation (MNS); electromyographic (EMG); flexor carpi radialis (FCR).

For each level of contraction (rest, 10%, 20% and 30% MVIC), three Mmax traces, twelve singlepulse MEPs (MEP_{test}) and twelve paired-pulse MEPs (MEP_{cond}) delivered with each of the TS intensities (120% and 130% of motor thresholds (MT)) were recorded per subject. The order of the different levels of contraction was randomized across participants. Single-pulse and paired-pulse TMS measures obtained with the two TS intensities performed in different recording blocks and the order of the blocks randomized across participants. The inter-trial intervals between single-pulses were at least 3 s [22]. During active conditions, the mean duration of the recording was set at 500 ms and participants were asked to relax after the stimulation for few seconds before the next trial and contracting again. This was done to avoid fatigue generated by holding the level of contraction across all consecutive trials.

2.3. EMG Recordings

After shaving and dry-cleaning the skin with alcohol, surface electromyographic (EMG) recordings of the right flexor carpi radialis (FCR) were made with two silver-chloride (Ag/AgCl) electrodes placed over the muscle belly with an interelectrode center-to-center distance of 2 cm, at 1/3 of the distance from the medial epicondyle of the humerus to the radial styloid process (Figure 1). A ground electrode was placed over the medial epicondyle to the radial styloid. The EMG signals were amplified (gain: 1000) and band-pass filtered (10–1000 Hz), digitized at a sampling rate of 2000 Hz and stored for off-line analysis (Biopac Systems Inc., Goleta, CA, USA).

2.4. Electrical Stimulation Protocol

To evoke Mmax, 1-ms square wave pulses were delivered over the median nerve near the cubital fossa via bipolar felt pad electrodes (8-mm diameter with 30-mm spacing) (Figure 1). The stimulation site, defined as the location that produced the largest Mmax, was first determined, then the electrodes, connected to a constant-current stimulator (DS7R, Digitimer, Welwyn Garden City, Hertfordshire, UK), were fixed at the arm by a Velcro strap. The intensity of stimulation was gradually increased until the amplitude of the M-wave did not increase any further. Then, 120% of this last intensity was used for Mmax recording, in order to ensure that the response lay in the plateau of its maximal value (mean intensity: 17.71 mA; range 13.2–28.2 mA).

2.5. Transcranial Magnetic Stimulation Protocol

Single-pulse and paired-pulse stimulations were applied with a 70-mm figure-of-eight coil connected to a monophasic Magstim BiStim² stimulator (The Magstim Co., Whitland, UK). The coil was placed over left M1, tangentially to the scalp with the handle pointing backward and laterally at 45° away from the midsagittal line, resulting in a postero-anterior current flow within M1. The optimal stimulation site on the scalp (hotspot) was defined as the location eliciting the largest MEP amplitude in the FCR muscle for a given intensity. This location was marked by a color marker on a tight-fitting cap worn by the participant. Because the aim of the current study was to compare the effect of different levels of muscle contraction (i.e., rest, 10, 20 and 30% of MVIC) and two TS intensities (i.e., 120% and 130% of the motor thresholds (MT)), MT were systematically calculated according to these different conditions. The resting motor threshold (rMT) was determined as the lowest stimulus intensity required to evoke at least 5 MEPs of 50 µV peak-to-peak amplitude out of 10 consecutive trials in the relaxed muscle [23]. For the 10%, 20% and 30% MVIC, the active motor threshold (aMT) was determined as the lowest stimulus intensity required to evoke at least 5 MEPs of 200 μ V peak-to-peak amplitude out of 10 consecutive trials [23]. Some participants (n = 4) presented an EMG level greater than 200 µV during muscle contraction. The aMT was then determined as the lowest stimulus intensity required to evoke at least 5 MEPs out of 10 consecutive trials with a peakto-peak amplitude greater than the 95% confidence interval of the prestimulus mean EMG activity for each contraction level [16]. For SICI measurements, the CS intensity was set at 70% rMT for rest conditions and 70% aMT for active conditions [9,16]. The inter-stimulus interval between TS and CS was set at 3 ms [5].

2.6. Data Analysis

Data analysis was performed using the Acqknowledge software package (Version 4.2, Biopac Systems Inc., Goleta, CA, USA).

The background root mean square of the surface EMG (EMG_{rms}) of the right FCR was calculated during the 100 ms epoch prior to the TMS artefact in order to quantify the pre-stimulus neuromuscular activation for all conditions. EMG_{rms} activity was normalized using the corresponding Mmax amplitude by the following ratio (EMG_{rms}/Mmax). EMG_{rms} activity was also normalized by using the maximal EMG_{rms} obtained during the participant's MVIC.

Responses were recorded in the right FCR muscle. The peak-to-peak amplitudes were measured for Mmax, MEP_{test} and MEP_{cond} in each experimental condition. Figure 2 shows typical raw Mmax

(2A), MEP_{test} (2B, white panel) and MEP_{cond} (2B, grey panel) elicited at 120% MT and MEP_{test} (2C, white panel) and MEP_{cond} (2C, grey panel) elicited at 130% MT.



Figure 2. Typical raw data recordings of the right FCR muscle depicted at rest, 10%, 20% and 30% of maximal voluntary isometric contraction (MVIC) for a representative participant. Black and grey vertical arrows indicate time of stimulation for test and conditioning stimuli, respectively. (**A**) Maximal M-wave (Mmax) induced by electrical stimulation. (**B**) Unconditioned (white) and conditioned (grey) motor-evoked potentials (MEPs) with the test stimulus induced at 120% of the motor threshold. (**C**) Unconditioned (white) and conditioned (grey) MEP with the test stimulus induced at 130% of the motor threshold. Note that the *y*-axis scale for MEPs amplitude differs between resting and active conditions (i.e., 0.2 mV vs. 1 mV) for a better graphical visualization.

In order to investigate how the estimation of SICI is influenced according to the normalization, SICI was expressed across all conditions using the following two equations:

(1) The 'usual' equation provides an estimation of SICI (SICI_{MEPtest}) by calculating the difference between the mean MEP_{cond} and the mean MEP_{test}, then expressed as a percentage of the mean MEP_{test} [9].

$$SICI (\%MEP_{test}) = \frac{MEP_{cond} - MEP_{test}}{MEP_{test}} \times 100$$
(1) (1)

(2) The second equation provides an estimation of SICI (SICI_{Mmax}) by calculating the difference between the mean MEP_{cond} and the mean MEP_{test} then expressed as a percentage of the mean Mmax [9,14].

SICI (%Mmax) =
$$\frac{\text{MEP}_{\text{cond}} - \text{MEP}_{\text{test}}}{\text{Mmax}} \times 100$$
 (2) (2)

For both equations, negative values indicate inhibition and positive values indicate facilitation.

2.7. Statistical Analysis

All statistical analyses were performed using IBM's SPSS software version 24 (SPSS Inc., Chicago, IL, USA). Normality of the data distributions was verified using the Shapiro–Wilk test and the assumption of normality was not violated for any of the data. Homogeneity of variances was assessed by Mauchly's test and a Greenhouse–Geiser correction was applied if the sphericity assumption was violated. Pre-planned post-hoc analyses were performed on significant interactions after applying a Bonferroni correction for multiple comparisons. Uncorrected degrees of freedom and corrected *p* values for multiple comparisons are reported in the results section. The α level for all analyses was fixed at 0.05. Partial eta squared (η_{p^2}) values are reported when results are statistically significant in order to express the portion of the total variance attributable to the tested factor or interaction. Values are reported as mean ± SD.

A first set of analyses were performed to control for potential methodological biases. The stimulation intensities (%MSO) for the MT as well as the EMG_{rms}/Mmax ratios and the Mmax amplitude obtained for each level of contraction were compared using a one-factor analyses of variance (ANOVA).

The normalized TMS data (MEP_{test}/Mmax; SICI_{MEPtest} and SICI_{Mmax}) were separately analysed with repeated-measure ANOVA with two within-subject factors: TS intensity₂ (120% rMT or aMT vs. 130% rMT or aMT) and levels of contraction₄ (i.e., rest, 10, 20 and 30% of MVIC).

In order to explore how SICI_{Mmax} can be predicted from the MEP_{test}/Mmax ratio, a regression coefficient analysis was estimated individually for each participant, according to procedures described by Pfister and colleagues [24]. Individual slopes, intercept, R² and *p* values were extracted and reported.

Then, to investigate how SICI_{MEPtest} and SICI_{Mmax} are influenced by the MEP_{test}/Mmax ratio that is, regardless of the level of contraction or the TS intensity, results were reorganised by pooling the MEP_{test}/Mmax ratio into 5% bins, which resulted in bins from 0–5% to 45–50% and then into 10% bins (50 to 60% and 60 to 70%) or a percentage corresponding to >70%. This also allowed to take into account inter-individual variability. Mean regression coefficients were calculated and reported for the two SICI normalizations.

3. Results

3.1. Methodological Considerations

The one-factor ANOVA conducted to compare the MT (rMT vs. aMT during 10% MVIC vs. 20% MVIC vs. 30% MVIC) showed a significant main effect ($F_{(3,42)} = 24.59$; p < 0.001; $\eta_p^2 = 0.637$, Table 1). Post-hoc comparisons revealed that only the rMT differs from all aMT (all p < 0.002). Post-hoc comparisons also indicate that, although systematically calculated for each level of contraction, the aMT did not significantly differ depending on the level of muscle contraction. An ANOVA performed on the mean Mmax amplitude obtained for each level of contraction showed no significant main effect ($F_{(3,42)} = 1.153$; p = 0.323, Table 1), indicating that the increase in the level of muscle contraction from rest to 30% MVIC did not affect Mmax amplitude.

Table 1. Comparison of motor thresholds (MT) (%maximum stimulator output (MSO)), single- and
paired-pulse TMS raw EMG $_{rms}$ for MEP data (μV), EMG $_{rms}$ expressed as a% of EMG $_{rms}$ obtained during
MVIC; Mmax amplitude (mV) and raw EMG _{ms} for Mmax data (μ V) across levels of muscle
contraction (%MVIC). Data represent mean ± SD. The MT for the Rest condition was significantly
higher than all the MT obtained for active conditions (10, 20 and 30% MVIC).

Variables		Levels of Muscle Contraction (%MVIC)				
		Rest	10%	20%	30%	
	MT (%MSO)	40.87 ± 7.3 **	31.73 ± 8.1	31.53 ± 7.2	31.27 ± 7	
	EMGrms MEP (µV)	1.97 ± 0.8	67.18 ± 28.2	113.58 ± 46.5	160.32 ± 64.5	
EMC	Grms (%EMGrms-MVIC)	0.31 ± 0.2	9.65 ± 5.7	16.06 ± 8.7	21.96 ± 10.5	
	Mmax (mV)	6.97 ± 3.7	6.82 ± 3.8	6.60 ± 3.7	6.70 ± 3.9	
E	EMGrms Mmax (µV)	1.75 ± 0.6	65.04 ± 27.7	114.83 ± 61	160.73 ± 78.8	

** *p* < 0.001. motor thresholds (MT); maximum stimulator output (MSO); root mean square (rms); electromyographic (EMG); motor-evoked potential (MEP); maximum voluntary isometric contraction (MVIC).

3.2. Neurophysiological Data

Figure 3 illustrates the EMG_{rms}/Mmax ratios pooled for single and paired pulses and the two TS intensities (120% and 130% MT). The ANOVA carried out on these EMG_{rms}/Mmax ratios revealed a significant main effect of the levels of contraction ($F_{(3,42)} = 83.32$; p < 0.001; $\eta_p^2 = 0.856$). Post hoc analyses showed a lower EMG_{rms}/Mmax ratio for the rest condition compared with the 10% MVIC (p < 0.001), 10% MVIC compared with 20% MVIC (p = 0.003) and 20% MVIC compared with the 30% MVIC (p = 0.001).





Figure 3. EMG_{rms}/Mmax ratios pooled for MEP_{test} (unconditioned) and MEP_{cond} (conditioned) and the two test stimulus (TS) intensities (120% and 130% of motor threshold (MT)). EMG_{rms} were recorded 100 ms before the delivery of each TMS pulse. ** p < 0.01

Figure 4A illustrates the MEP_{test}/Mmax ratios according to the TS intensity and the levels of contraction. The ANOVA revealed a main effect of the TS intensity ($F_{(1,14)}$ = 26.97; *p* < 0.001; η_p^2 = 0.658) showing that the MEP_{test}/Mmax ratio was greater with a TS intensity set at 130% MT than with a TS intensity set at 120% MT. A significant large effect of the levels of muscle contraction was also observed ($F_{(3,42)}$ = 120.01; *p* < 0.001; η_p^2 = 0.896). Post hoc analyses showed a significant increase in the MEP_{test}/Mmax for 10% MVIC compared with rest (*p* < 0.001), for 20% MVIC compared with 10% MVIC (*p* < 0.001) and for 30% MVIC compared with 20% MVIC (*p* < 0.001). The TS intensity by levels of contraction interaction was not significant ($F_{(3,42)}$ = 2.16; *p* = 0.107).

Figure 4B illustrates the SICI when normalized to MEP_{test}. The SICI_{MEPtest} did not differ according to the TS intensity ($F_{(1,14)} < 1$; p = 0.538) but the levels of muscle contraction main effect was significant

 $(F_{(3,42)} = 5.15; p = 0.020; \eta_p^2 = 0.268)$. Post-hoc analyses showed that SICI_{MEPtest} was higher when measured at rest than during 30% MVIC (p = 0.043). The TS intensity by levels of muscle contraction interaction was not significant ($F_{(3,42)} = 1.99; p = 0.161$).



Figure 4. (**A**) MEP_{test}/Mmax ratios according to the TS intensity and the levels of contraction. (**B**) Shortinterval intracortical inhibition (SICI) expressed as percentage of MEP_{test} (SICI_{MEPtest}) according to the TS intensity and the levels of contraction. (**C**) SICI expressed in percentage of Mmax (SICI_{Mmax}) according to the TS intensity and the levels of contraction. * p < 0.05; ** p < 0.001.

Figure 4C illustrates the SICI when normalized to Mmax. Similar to the ratio MEP_{test}/Mmax, the SICI_{Mmax} was significantly higher with a TS intensity of 130% MT when compared with a TS intensity of 120% MT ($F_{(1,14)} = 6.73$; p = 0.021; $\eta_p^2 = 0.325$). A significant large main effect of the levels of contraction was found ($F_{(3,42)} = 63.94$; p < 0.001; $\eta_p^2 = 0.830$). Post-hoc analyses revealed that SICI_{Mmax} was higher at 10% MVIC when compared with rest (p < 0.001) and higher at 20% MVIC when compared with 10% MVIC (p = 0.006). The TS intensity by levels of muscle contraction interaction failed to reach significance ($F_{(3,42)} = 1.73$; p = 0.199).

As reported in Table 2, the regression analysis was also significant for all subjects and provided evidence for a strong linear relationship between the SICI_{Mmax} estimation and the MEP_{test}/Mmax ratios. This indicates that it is possible to predict SICI_{Mmax} in an individual from a specific MEP_{test}/Mmax ratio.

Subject	Slope	Intercept	R ²	<i>p</i> Value	Range of the MEP _{test} /Mmax Ratios (%)
1	-0.553	0.118	0.911	< 0.001	1.19–46.9
2	-0.798	0.508	0.944	< 0.001	2.97-32.91
3	-0.593	-0.320	0.879	= 0.001	7.54-46.85
4	-0.546	-1.474	0.895	< 0.001	1.23-46.74
5	-0.518	-1.143	0.806	= 0.002	6.40-45.38
6	-0.693	0.728	0.941	< 0.001	1.11-47.68
7	-0.725	-1.884	0.943	< 0.001	6.07-72.37
8	-0.455	-6.172	0.848	= 0.001	10.66–74.25
9	-0.626	-1.282	0.898	< 0.001	2.44-69.65
10	-0.486	-3.472	0.947	< 0.001	8.28-62.33
11	-0.610	0.231	0.961	< 0.001	1.52-58.23
12	-0.466	-2.872	0.856	= 0.001	5.38-49.67
13	-0.405	-5.942	0.791	= 0.003	7.29–66.39
14	-0.385	-1.949	0.921	< 0.001	1.75–79.66
15	-0.671	-1.563	0.937	< 0.001	5.84-89.71
Mean	-0.569	-1.706	0.898		4.65–59.25
SD	0.120	2	0.053		3.09–15.78

Table 2. Regression analysis between MEP_{test}/Mmax ratio (independent variable) and SICI_{Mmax} (dependent variable) estimated individually for each subject. Individual slopes, intercept, R², *p* values and range of the MEP/Mmax ratios were extracted and reported.

SICI_{MEPtest} (Figure 5A) and SICI_{Mmax} (Figure 5B) plotted according to the MEP_{test}/Mmax values pooled into different bins. First, a simple linear regression was calculated on mean bins values in order to predict the SICI_{MEPtest} based on the MEP_{test}/Mmax ratio (Figure 5C). A non-significant regression equation was found ($F_{(1,12)} < 1$, p = 0.369), with a low R² of 0.081. This indicates that MEP_{test}/Mmax ratio cannot predict the SICI_{MEPtest}. Then, a simple regression was calculated on mean values to predict the SICI_{Mmax} based on the MEP_{test}/Mmax ratio (Figure 5D). A significant regression equation was found ($F_{(1,12)} = 553.2$, p < 0.001), with a large R² of 0.980. Crucially, this indicates that the relationship between SICI and corticospinal excitability is relative to/dependent on the proportions of the spinal motoneurones pool activated by TMS, no matter how the corticospinal excitability is actively (levels of contraction) or passively (TMS intensity) increased.



Figure 5. SICI normalized to MEP_{test} (SICI_{MEPtest}) (**A**) and SICI normalized to Mmax (SICI_{Mmax}) (**B**) organized into different bins based on the MEP_{test}/Mmax ratio. Grey bars represent the mean value for each bin, with standard error of the mean. Black dots represent individual values included in each bin. Non-significant regression plots between MEP_{test}/Mmax ratio SICI_{MEPtest} (**C**) and significant regression plots between MEP_{test}/Mmax ratio SICI_{MEPtest} (**C**) and significant regression plots between MEP_{test}/Mmax ratio SICI_{MEPtest} (**C**) and significant regression plots between MEP_{test}/Mmax ratio SICI_{Mmax} (**D**) Black dots in regressions represent the mean value for each bin. Grey lines in the regression figures represent 95% confident interval.

4. Discussion

This study investigated the modulation of SICI when normalizing or not with the proportion of spinal motoneurones activated (SICI_{Mmax} vs. SICI_{MEPtest}), with varying MEP_{test} amplitude. Our findings highlight the dependence of SICI measurement upon the proportion of motoneurones activated, i.e., SICI_{Mmax}, regardless of the muscle state and TS intensity. The strong linear relationship between SICI_{Mmax} estimation and MEP_{test}/Mmax ratio provides a relevant predictor of SICI for each individual. The main findings and their implications for future studies are discussed below.

4.1. Modulation of Corticospinal Excitability and SICIMEPtest

In the current study, we first measured the amplitude of unconditioned MEP (MEP_{test}) by handling muscle state (at rest or at three levels of contraction) and TS intensity (120 or 130% of MT for each level of muscle contraction). As expected, corticospinal excitability increased with increasing muscle contraction and TS intensity [25,26].

When normalizing SICI with MEP_{test} (SICI_{MEPtest}), we did not find any SICI_{MEPtest} modulation according to TS intensity. The current findings only revealed one significant difference between muscle states, with a reduction in SICI_{MEPtest} when tested at 30%MVIC and compared with rest. These results are therefore similar to those obtained in studies that evaluate SICI with no account of the proportion of motoneurones activated by the TS. Indeed, it is known that voluntary muscle contraction reduced SICI_{MEPtest} [16,27], particularly when tested with high levels of muscle contraction (>40% MVIC) [17,19]. One explanation is that during high levels of voluntary muscle contraction, the excitability of the spinal motoneurones pool is increased and the later indirect waves (called I3) generated by the TS (fixed to induce 1 mV MEP amplitude in the majority of previous studies) no longer contribute to the production of the MEPtest [28]. Indeed, the EMG response elicited by TS is

usually composed of different waves, namely D-waves and I-waves. While direct D-wave is the first latency response originating from the activation of the pyramidal tract axons, the later indirect I-waves, numbering three waves (I1, I2, I3), are thought to result from the pyramidal neuron activation [29]. Because the CS predominantly inhibits the I3 [30], a decrease in SICI_{MEPtest} is observed at higher contraction levels (when there is no I3 to suppress) [18,27,31]. Finally, one has to keep in mind that a 1 mV fixed MEP amplitude across rest and active conditions (1) represents a different proportion of the spinal motoneurones pool activated by adjusted TMS intensity within an individual, and (2) does not result from the same contribution of descending waves activated by TMS, whereas SICI acts selectively on some of these descending waves.

Importantly, with low levels of muscle contraction [17,19] or different MEP_{test} amplitudes evoked at 120 or 130% of MT [11,32], the amount of SICI_{MEPtest} remains unchanged. As mentioned by Lackmy and Marchand-Pauvert (2010), this result seems logical from a purely mathematical point of view given the linear relationship between the MEP_{cond} and MEP_{test} amplitudes [9]. Therefore, the MEP_{cond}-MEP_{test}/MEP_{test} or MEP_{cond}/MEP_{test} ratios result in a slope of an approximately linear relationship. This mathematical relationship explains why the SICI_{MEPtest} is not influenced by the MEP_{test} amplitude, whether the latter is increased by the level of muscle contraction or by TS intensity.

4.2. Modulation of SICI_{Mmax} and Its Relationship with MEP_{test}/Mmax Ratio

By taking into account the proportion of motoneurones activated by the TS to quantify SICI, we found more inhibition with higher MEP_{test}/Mmax ratio, both at rest and during the active muscle conditions. A major finding of the current study is the strong individual relationship between the SICI_{Mmax} estimation and the MEP_{test}/Mmax ratios, which is ultimately independent from experimental conditions (muscle state and TS intensity). Increasing the level of muscle contraction or the TS intensity both led to a greater proportion of large spinal motoneurones activated by TMS and an increase in the MEP_{test} amplitude. Because of their higher threshold (when compared with smaller low-threshold motoneurones), these large motoneurones are more sensitive to a decrease in corticospinal input, induced by the CS [33]. By considering the recruitment order of motoneurones, it is possible that SICI_{Mmax} had a greater effect on the larger MEP_{test} amplitude and therefore the MEP_{test}/Mmax ratio. Therefore, the greater the MEP_{test}/Mmax ratio, the greater the SICI_{Mmax}.

The individual relationship obtained makes it possible to individually predict SICI_{Mmax} from a specific MEP_{test}/Mmax ratio. This result is of particular methodological importance given the crucial contribution of SICI modulations to motor skill acquisition [34,35] or acute and multi-session strength training programs [3,36,37]. Several recent articles highlighted the high heterogeneity in the setting parameters, such as muscle state, used to evaluate SICI modulation before and after motor practice [2,17,38,39]. Some authors argued that, when the muscle is active, measurements of intracortical and corticospinal excitability are more sensitive and specific to motor learning than the same tests performed at rest after motor learning/training [40]. This statement, however, does not indicate the level(s) of muscle contraction to choose during testing. The findings of the current study are also interesting for protocols assessing SICI changes following an acute fatiguing exercise [40]. In these protocols, post-test measures must be performed immediately following the exercise in order to avoid a recovery influence. Fatigue is known to increase the size of MEP_{test} [41,42]. Therefore, adjusting TS intensity at the beginning of post-test SICI measurement to obtain a similar MEP_{test} amplitude than during pre-test (such as 1 mV) is a time-consuming procedure that could bias the interpretation of fatigue-induced modulation of SICI.

It should be noted that while we observed a greater amount of SICI_{Mmax} with a greater MEP_{test}/Mmax ratio, Lackmy and Marchand-Pauvert (2010) reported a decrease in SICI_{Mmax} when the MEP_{test}/Mmax ratio was higher than 30%. In their study, TMS-evoked responses were recorded in small hand muscles with increasing TS intensities for a given CS. The authors found a peak of inhibition when the MEP_{test}/Mmax ratio was approximately equal to 20–30%, explained by a greater inhibitory effect of SICI on large and high-threshold motoneurones (i.e., larger MEP_{test}/Mmax ratio). This peak of inhibition was then followed by a decrease in the SICI_{Mmax}. When increasing TS intensities, and therefore MEP_{test}/Mmax ratios, greater short-latency direct-waves (D-waves) are

recruited [43]. It is known that these D-waves are less sensitive to SICI [8,14,30], resulting in SICI_{Mmax} decrease with increasing MEP_{test}/Mmax ratios. Concurrently, in this study, SICI_{Mmax}, MEP_{test} and Mmax were tested in a more proximal muscular group (i.e., forearm muscle) than the above-cited study. This choice of testing FCR muscle was mainly motivated by the common use of wrist muscles in numerous strength-training programs that require pre-intervention and post-intervention SICI measurements [37,38,44–46]. The recruitment and discharge properties of motoneurones are known to be different between small hand muscles compared with more proximal arm muscles [47]. During voluntary contractions, MEPs amplitude evoked in a hand muscle at a fixed TS intensity was almost saturated at 20%MVIC whereas MEPs amplitude evoked at the same intensity in the FCR reached a plateau at 40%MVIC [47]. These characteristics probably explain why, in the current study, we did not observe a decrease in SICI_{Mmax} with MEP_{test}/Mmax ratio greater than 30–40%.

4.3. Methodological Recommendations

By calculating the individual slopes, intercept, R² and *p*-values between the MEPtest/Mmax ratio and the SICI_{Mmax} before an intervention (as carried out in the present study), it becomes possible to overcome the testing of a specific level of muscle contraction, or to perform a size MEPtest adjustment in order to test SICIMmax modulation after motor practice in healthy subjects. It only requires the calculation of the mean SICIMmax for a specific MEPtest/Mmax ratio after any intervention. Then, it is possible to compare this SICIMmax value with the equation of the regression obtained before the intervention in order to evaluate the SICI_{Mmax} modulation induced. There are two possible results: (1) the value obtained in post-tests is aligned with the theoretical linear relationship found in pre-tests, indicating no change in SICI_{Mmax} following the intervention; and (2) the value obtained in post-tests is above/under the theoretical line, indicating that for a given MEPtest/Mmax ratio, SICIMmax decreases/increases following the intervention. Importantly, in the current study, the individual regression analyses are based on eight data points per individual (i.e., 2 TS intensities and four levels of muscle contractions). A quicker approach can be considered, with a more restricted number of points. For example, individual regression analyses based on only four points, corresponding to different levels of muscle contractions with a unique TS intensity, also lead to strong individual relationships between the SICIMmax estimation and the MEPtest/Mmax ratios (see Table S1). It is important to keep in mind that these methodological recommendations are based on results obtained in a study completed acutely. Since Mmax does not always remain constant during the course of an experiment [48] and has been shown to be altered following strength-training [49], it is crucial to repeat the testing for each experimental condition, including during post-test measurements.

4.4. Limitations

The current study presents certain limitations that should be noted. First, the number of pulses selected to evaluate SICI is quite smaller than the one recommended for highly reliable SICI estimates (i.e., more than 25 pulses) [50]. This choice was made to avoid any fatigue effect that might have potentially confounded the results obtained. Moreover, both SICIMEPTEST and SICIMERA modulations also depend on CS intensity and interstimulus interval (ISI) duration [6,12,51-53], two fixed parameters whose effects were not assessed in the current study. The CS intensity was fixed at 70% of MT calculated according to each levels of contraction. This was done based on a previous study showing that during active condition, the higher level of inhibition was obtained delivering a CS of 70% aMT [16]. Of note, a recent study has demonstrated that the CS intensity (60%, 70% or 80% aMT) did not alter the modulation of SICIMEPTest assessed at different muscle contraction levels [17]. Concerning the ISI, it is known that SICIMEPTEST consists of two phases of inhibition according to the ISI duration [6,12]. The first one is observed at ISI = 1 ms and is thought to reflect the refractoriness of axons of cortical interneurons subliminally activated by a subthreshold CS [6], and/or extrasynaptic GABA tone [54]. The second one observed at ISI = 2.5/3 ms accounted for synaptic inhibition mediated by the GABAA receptors [6,12,55]. The 3-ms ISI was selected in the current study as this duration consistently demonstrates a clear peak of inhibition [53,56]. However, future studies should address whether CS intensity and ISI duration can affect SICI_{Mmax} modulation. Finally, only one forearm muscle was tested,

and we did not consider (1) antagonist muscle and (2) synergistic muscles that are also relevant to force production without a concomitant increase in the EMG of the FCR agonist muscle. At this stage, the results cannot be generalized to all muscle groups and deserve further investigations.

5. Conclusions

The current study provides new methodological insights into how SICI, representing an inhibitory phenomenon known to act purely at the cortical level, is ultimately dependent upon the proportion of motoneurones activated by TMS. Moreover, based on the results obtained, our study suggests a new way to evaluate SICI_{Mmax} modulation that makes it possible to get away from important methodological constraints related to pre/post intervention measures. This may have practical implications for the assessment of cortical adaptations that contribute to functional changes induced by motor practice, motor learning, motor exercise or strength training in future studies.

Supplementary Materials: The following are available online at www.mdpi.com/2076-3425/10/7/433/s1, Table S1: Regression analysis between MEP_{test}/Mmax ratio (independent variable) and SICI_{Mmax} (dependent variable) estimated individually for each subject with TS intensity set at 120%MT (left) and 130%MT (right).

Author Contributions: Conceptualization, C.N., S.G., A.M. and F.L.; methodology, C.N., S.G., A.M. and F.L.; analysis, C.N., S.G., A.M. and F.L.; investigation, C.N.; writing—original draft preparation, C.N., S.G., A.M. and F.L.; writing—review and editing, C.N., S.G., A.M. and F.L.; supervision, F.L.; funding acquisition, F.L. All authors have read and agreed to the published version of the manuscript.

Funding: This work was funded by the French "Investissements d'Avenir" program, project ISITE-BFC (contract ANR-15-IDEX-0003).

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Data Available: https://osf.io/jhaus/.

References

- Reis, J.; Swayne, O.B.; Vandermeeren, Y.; Camus, M.; Dimyan, M.A.; Harris-Love, M.; Perez, M.A.; Ragert, P.; Rothwell, J.C.; Cohen, L.G. Contribution of transcranial magnetic stimulation to the understanding of cortical mechanisms involved in motor control. *J. Physiol.* 2008, 586, 325–351.
- 2. Kidgell, D.J.; Bonanno, D.R.; Frazer, A.K.; Howatson, G.; Pearce, A.J. Corticospinal responses following strength training: A systematic review and meta-analysis. *Eur. J. Neurosci.* **2017**, *46*, 2648–2661.
- 3. Weier, A.T.; Pearce, A.J.; Kidgell, D.J. Strength training reduces intracortical inhibition. *Acta Physiol.* **2012**, 206, 109–119.
- 4. Siddique, U.; Rahman, S.; Frazer, A.; Leung, M.; Pearce, A.J.; Kidgell, D.J. Task-dependent modulation of corticospinal excitability and inhibition following strength training. *J. Electromyogr. Kinesiol.* **2020**, *52*, 102411.
- 5. Kujirai, T.; Caramia, M.D.; Rothwell, J.C.; Day, B.L.; Thompson, P.D.; Ferbert, A.; Wroe, S.; Asselman, P.; Marsden, C.D. Corticocortical inhibition in human motor cortex. *J. Physiol.* **1993**, *471*, 501–519.
- 6. Fisher, R.J.; Nakamura, Y.; Bestmann, S.; Rothwell, J.C.; Bostock, H. Two phases of intracortical inhibition revealed by transcranial magnetic threshold tracking. *Exp. Brain Res.* **2002**, *143*, 240–248.
- 7. Di Lazzaro, V.; Restuccia, D.; Oliviero, A.; Profice, P.; Ferrara, L.; Insola, A.; Mazzone, P.; Tonali, P.; Rothwell, J.C. Magnetic transcranial stimulation at intensities below active motor threshold activates intracortical inhibitory circuits. *Exp. Brain Res.* **1998**, *119*, 265–268.
- 8. Di Lazzaro, V.; Rothwell, J.; Capogna, M. Noninvasive stimulation of the human brain: Activation of multiple cortical circuits. *Neuroscientist* **2018**, *24*, 246–260.
- 9. Lackmy, A.; Marchand-Pauvert, V. The estimation of short intra-cortical inhibition depends on the proportion of spinal motoneurones activated by corticospinal inputs. *Clin. Neurophysiol.* **2010**, *121*, 612–621.
- 10. Garry, M.I.; Thomson, R.H.S. The effect of test TMS intensity on short-interval intracortical inhibition in different excitability states. *Exp. Brain Res.* **2009**, *193*, 267–274.

- 11. Sanger, T.D.; Garg, R.R.; Chen, R. Interactions between two different inhibitory systems in the human motor cortex. *J. Physiol.* **2001**, *530*, 307–317.
- 12. Roshan, L.; Paradiso, G.O.; Chen, R. Two phases of short-interval intracortical inhibition. *Exp. Brain Res.* **2003**, *151*, 330–337.
- 13. Grosprêtre, S.; Papaxanthis, C.; Martin, A. Modulation of spinal excitability by a sub-threshold stimulation of M1 area during muscle lengthening. *Neuroscience* **2014**, *263*, 60–71.
- 14. Opie, G.M.; Semmler, J.G. Modulation of short- and long-interval intracortical inhibition with increasing motor evoked potential amplitude in a human hand muscle. *Clin. Neurophysiol.* **2014**, *125*, 1440–1450.
- 15. Ridding, M.C.; Taylor, J.L.; Rothwell, J.C. The effect of voluntary contraction on cortico-cortical inhibition in human motor cortex. *J. Physiol.* **1995**, *487*, 541–548.
- 16. Ortu, E.; Deriu, F.; Suppa, A.; Tolu, E.; Rothwell, J.C. Effects of volitional contraction on intracortical inhibition and facilitation in the human motor cortex. *J. Physiol.* **2008**, *586*, 5147–5159.
- 17. Hendy, A.M.; Ekblom, M.M.; Latella, C.; Teo, W.P. Investigating the effects of muscle contraction and conditioning stimulus intensity on short-interval intracortical inhibition. *Eur. J. Neurosci.* **2019**, *50*, 3133–3140.
- 18. Zoghi, M.; Nordstrom, M.A. Progressive suppression of intracortical inhibition during graded isometric contraction of a hand muscle is not influenced by hand preference. *Exp. Brain Res.* **2007**, 177, 266–274.
- 19. Rantalainen, T.; Weier, A.; Leung, M.; Brandner, C.; Spittle, M.; Kidgell, D. Short-interval intracortical inhibition is not affected by varying visual feedback in an isometric task in biceps brachii muscle. *Front. Hum. Neurosci.* **2013**, *7*, 68.
- 20. Oldfield, R.C. The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia* **1971**, *9*, 97–113.
- 21. Rossi, S.; Hallett, M.; Rossini, P.M.; Pascual-Leone, A. Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clin. Neurophysiol.* **2009**, *120*, 2008–2039.
- 22. Rothwell, J.C.; Hallett, M.; Berardelli, A.; Eisen, A.; Rossini, P.; Paulus, W. Magnetic stimulation: Motor evoked potentials. *Electroencephalogr. Clin. Neurophysiol. Suppl.* **1999**, *52*, 97–103.
- 23. Rossini, P.; Barker, A.; Berardelli, A. Non-invasive electrical and magnetic stimulation of the brain, spinal cord and roots: Basic principles and procedures for routine clinical application. Report of an IFCN. *Electroencephalogr. Clin. Neurophysiol.* **1994**, *91*, 79–92.
- 24. Pfister, R.; Schwarz, K.; Carson, R.; Jancyzk, M. Easy methods for extracting individual regression slopes: Comparing SPSS, R, and Excel. *Tutor. Quant. Methods Psychol.* **2013**, *9*, 72–78.
- 25. Valls-Solé, J.; Pascual-Leone, A.; Brasil-Neto, J.P.; Cammarota, A.; McShane, A.; Hallett, M. Abnormal facilitation of the response to transcranial magnetic stimulation in patients with parkinson's disease. *Neurology* **1994**, *44*, 735–741.
- 26. van den Bos, M.A.J.; Geevasinga, N.; Menon, P.; Burke, D.; Kiernan, M.C.; Vucic, S. Physiological processes influencing motor-evoked potential duration with voluntary contraction. *J. Neurophysiol.* **2017**, *117*, 1156–1162.
- 27. Zoghi, M.; Pearce, S.L.; Nordstrom, M.A. Differential modulation of intracortical inhibition in human motor cortex during selective activation of an intrinsic hand muscle. *J. Physiol.* **2003**, *550*, 933–946.
- 28. Di Lazzaro, V.; Restuccia, D.; Oliviero, A.; Profice, P.; Ferrara, L.; Insola, A.; Mazzone, P.; Tonali, P.; Rothwell, J.C. Effects of voluntary contraction on descending volleys evoked by transcranial stimulation in conscious humans. *J. Physiol.* **1998**, *508*, 625–633.
- 29. Terao, Y.; Ugawa, Y. Basic Mechanisms of TMS. J. Clin. Neurophysiol. 2002, 19, 322-343.
- 30. Di Lazzaro, V.; Profice, P.; Ranieri, F.; Capone, F.; Dileone, M.; Oliviero, A.; Pilato, F. I-wave origin and modulation. *Brain Stimul.* **2012**, *5*, 512–525.
- 31. Hanajima, R.; Ugawa, Y.; Terao, Y.; Sakai, K.; Furubayashi, T.; Machii, K.; Kanazawa, I. Paired-pulse magnetic stimulation of the human motor cortex: Differences among I waves. *J. Physiol.* **1998**, *509*, 607–618.
- 32. Wagle-Shukla, A.; Ni, Z.; Gunraj, C.A.; Bahl, N.; Chen, R. Effects of short interval intracortical inhibition and intracortical facilitation on short interval intracortical facilitation in human primary motor cortex. *J. Physiol.* **2009**, *587*, 5665–5678.
- 33. Lackmy-Vallee, A.; Giboin, L.S.; Marchand-Pauvert, V. Non-linear input-output properties of the cortical networks mediating TMS-induced short-interval intracortical inhibition in humans. *Eur. J. Neurosci.* **2012**, *35*, 457–467.

- 34. Liepert, J.; Classen, J.; Cohen, L.G.; Hallett, M. Task-dependent changes of intracortical inhibition. *Exp. Brain Res.* **1998**, *118*, 421–426.
- 35. Perez, M.A.; Lungholt, B.K.S.; Nyborg, K.; Nielsen, J.B. Motor skill training induces changes in the excitability of the leg cortical area in healthy humans. *Exp. Brain Res.* **2004**, *159*, 197–205.
- Goodwill, A.M.; Pearce, A.J.; Kidgell, D.J. Corticomotor plasticity following unilateral strength training. *Muscle Nerve* 2012, 46, 384–393.
- 37. Hendy, A.M.; Kidgell, D.J. Anodal tDCS applied during strength training enhances motor cortical plasticity. *Med. Sci. Sports Exerc.* **2013**, *45*, 1721–1729.
- Mason, J.; Frazer, A.K.; Pearce, A.J.; Goodwill, A.M.; Howatson, G.; Jaberzadeh, S.; Kidgell, D.J. Determining the early corticospinal-motoneuronal responses to strength training: A systematic review and meta-analysis. *Rev. Neurosci.* 2019, 30, 463–476.
- 39. Berghuis, K.M.M.; Veldman, M.P.; Solnik, S.; Koch, G.; Zijdewind, I.; Hortobágyi, T. Neuronal mechanisms of motor learning and motor memory consolidation in healthy old adults. *Age* **2015**, *37*, 53.
- 40. Gruet, M.; Temesi, J.; Rupp, T.; Levy, P.; Millet, G.Y.; Verges, S. Stimulation of the motor cortex and corticospinal tract to assess human muscle fatigue. *Neuroscience* **2013**, *231*, *384–399*.
- 41. Latella, C.; Hendy, A.; Vanderwesthuizen, D.; Teo, W.-P. The modulation of corticospinal excitability and inhibition following acute resistance exercise in males and females. *Eur. J. Sport Sci.* **2018**, *18*, 984–993.
- 42. Latella, C.; Teo, W.-P.; Harris, D.; Major, B.; VanderWesthuizen, D.; Hendy, A.M. Effects of acute resistance training modality on corticospinal excitability, intra-cortical and neuromuscular responses. *Eur. J. Appl. Physiol.* **2017**, *117*, 2211–2224.
- 43. Di Lazzaro, V.; Ziemann, U.; Lemon, R.N. State of the art: Physiology of transcranial motor cortex stimulation. *Brain Stimul.* **2008**, *1*, 345–362.
- Mason, J.; Howatson, G.; Frazer, A.K.; Pearce, A.J.; Jaberzadeh, S.; Avela, J.; Kidgell, D.J. Modulation of intracortical inhibition and excitation in agonist and antagonist muscles following acute strength training. *Eur. J. Appl. Physiol.* 2019, 119, 2185–2199.
- 45. Nuzzo, J.L.; Barry, B.K.; Jones, M.D.; Gandevia, S.C.; Taylor, J.L. Effects of Four Weeks of Strength Training on the Corticomotoneuronal Pathway. *Med. Sci. Sports Exerc.* **2017**, *49*, 2286–2296.
- 46. Mason, J.; Frazer, A.; Horvath, D.M.; Pearce, A.J.; Avela, J.; Howatson, G.; Kidgell, D. Adaptations in corticospinal excitability and inhibition are not spatially confined to the agonist muscle following strength training. *Eur. J. Appl. Physiol.* **2017**, *117*, 1359–1371.
- Wu, L.; Goto, Y.; Taniwaki, T.; Kinukawa, N.; Tobimatsu, S. Different patterns of excitation and inhibition of the small hand and forearm muscles from magnetic brain stimulation in humans. *Clin. Neurophysiol.* 2002, 113, 1286–1294.
- 48. Crone, C.; Johnsen, L.L.; Hultborn, H.; Orsnes, G.B. Amplitude of the maximum motor response (Mmax) in human muscles typically decreases during the course of an experiment. *Exp. Brain Res.* **1999**, *124*, 265–270.
- 49. Nuzzo, J.L.; Barry, B.K.; Gandevia, S.C.; Taylor, J.L. Acute Strength Training Increases Responses to Stimulation of Corticospinal Axons. *Med. Sci. Sports Exerc.* **2016**, *48*, 139–150.
- 50. Biabani, M.; Farrell, M.; Zoghi, M.; Egan, G.; Jaberzadeh, S. The minimal number of TMS trials required for the reliable assessment of corticospinal excitability, short interval intracortical inhibition, and intracortical facilitation. *Neurosci. Lett.* **2018**, *674*, 94–100.
- 51. Ibáñez, J.; Spampinato, D.A.; Paraneetharan, V.; Rothwell, J.C. SICI during changing brain states: Differences in methodology can lead to different conclusions. *Brain Stimul.* **2020**, *13*, 353–356.
- 52. Neige, C.; Rannaud Monany, D.; Stinear, C.M.; Byblow, W.D.; Papaxanthis, C.; Lebon, F. Unravelling the Modulation of Intracortical Inhibition During Motor Imagery: An Adaptive Threshold-Hunting Study. *Neuroscience* **2020**, *434*, 102–110.
- 53. Vucic, S.; Cheah, B.C.; Krishnan, A.V.; Burke, D.; Kiernan, M.C. The effects of alterations in conditioning stimulus intensity on short interval intracortical inhibition. *Brain Res.* **2009**, *1273*, 39–47.
- 54. Stagg, C.J.; Bestmann, S.; Constantinescu, A.O.; Moreno Moreno, L.; Allman, C.; Mekle, R.; Woolrich, M.; Near, J.; Johansen-Berg, H.; Rothwell, J.C. Relationship between physiological measures of excitability and levels of glutamate and GABA in the human motor cortex. *J. Physiol.* **2011**, *589*, 5845–5855.

- 55. Peurala, S.H.; Müller-Dahlhaus, J.F.M.; Arai, N.; Ziemann, U. Interference of short-interval intracortical inhibition (SICI) and short-interval intracortical facilitation (SICF). *Clin. Neurophysiol.* **2008**, *119*, 2291–229.
- 56. Cirillo, J.; Byblow, W.D. Threshold tracking primary motor cortex inhibition: The influence of current direction. *Eur. J. Neurosci.* **2016**, *44*, 2614–2621.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).