



**HAL**  
open science

# Degeneracy of algebro geometric solutions of the Toda equation and Sato formulas

Pierre Gaillard

► **To cite this version:**

Pierre Gaillard. Degeneracy of algebro geometric solutions of the Toda equation and Sato formulas. 2024. hal-04769964

**HAL Id: hal-04769964**

**<https://u-bourgogne.hal.science/hal-04769964v1>**

Preprint submitted on 6 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Degeneracy of algebro geometric solutions of the Toda equation and Sato formulas

<sup>+</sup>Pierre Gaillard,

<sup>+</sup> Université de Bourgogne Franche-Comté, Dijon, France :  
e-mail: Pierre.Gaillard@u-bourgogne.fr,

## Abstract

We know that the degeneracy of solutions to PDE's given in terms of theta functions on Riemann surfaces gives important results about particular solutions as in the case of the NLS equation.

Here we degenerate the so called finite gap solutions of the Toda lattice equation from the general formulation in terms of abelian functions when the gaps tends to points.

This degeneracy allows to recover the Sato formulas, without using inverse scattering theory or geometric or representation theoretic methods.

## MSC2020 numbers:

35 – *XX*, 35*Cxx*, 35*Qxx*

## 1 Introduction

Different formulations can be chosen to represent the Toda equation. In original work of 1967 [93], the vibration of a uniform chain of particles with nonlinear interaction was studied, and Toda considered the following equation for the  $n$ -th particle in the chain

$$m\partial_x^2(u_n) = -\partial_r(\phi)(u_n - u_{n-1}) + \partial_r(\phi)(u_{n+1} - u_n), \quad (1)$$

where  $m$  stands for the mass of the particles,  $\phi(r)$  the interaction energy between adjacent particles. Some solutions were constructed for particular interactions  $\phi$ .

The same equation was considered with  $m = 1$  by Date and Tanaka in 1976 [3] and they constructed some general solutions but in terms of integrals difficult to use.

In the appendix of the paper of Dubrovin [6] published in 1981, Krichever considered the non abelian version of this equation and gave solutions in terms of theta function on Riemann surfaces.

Matveev and Stahlhoffen [91] considered in 1995 the following version of the Toda equation

$$\partial_x^2(u_n) = \exp(u_{n-1} - u_n) - \exp(u_n - u_{n+1}). \quad (2)$$

They used the Darboux transformation to construct solutions of this equation in terms of Casoratians.

In 2014, Zhang and Zhou [95] used the following representation of the Toda equation

$$\partial_x \partial_t(u_n) = (\partial_t(u_n) + \alpha(t))(u_{n-1} - 2u_n + u_{n+1}). \quad (3)$$

They used the generalization of the exp-function to construct multiwaves solutions of this equation.

More recently, Sun, Ma and Yu [95] were interested in 2020 in the following representation of the Toda equation

$$\Delta(u_n) = 4(\exp(u_{n-1} - u_n) - \exp(u_n - u_{n+1})). \quad (4)$$

They used a logarithmic transformation to get some particular solutions in terms of logarithms.

Duarte [4] considered in 2020 the following representation of the Toda equation

$$\begin{aligned} & \alpha \partial_x^2(u(x, y, n)) + \beta \partial_y^2(u(x, y, n)) \\ & = \exp(u(x, y, n-1) - u(x, y, n)) - \exp(u(x, y, n) - u(x, y, n+1)). \end{aligned}$$

He used a particular ansatz combined with the properties of Laplace's equation to construct some solutions in terms of trigonometric functions.

Schiebold and Nilson [88] studied in 2020 another version of the Toda equation in the form

$$\partial_{xy}^2(\log(1 + u_n)) = u_{n+1} - 2u_n + u_{n-1}. \quad (5)$$

They constructed solutions in the frame of linear algebra by means of determinants.

Here, we consider the Toda equation in the following representation [1, 5, 81, 90]

$$\begin{cases} \frac{r_n}{r_{n+1}} \sqrt{c_n} \psi_{n+1} + v_n \psi_n + \frac{r_n}{r_{n-1}} \sqrt{c_{n-1}} \psi_{n-1} = \lambda \psi_n \\ \partial_t \psi_n = \frac{r_n}{2r_{n+1}} \sqrt{c_n} \psi_{n+1} + \partial_t(\ln r_n) \psi_n - \frac{r_n}{2r_{n-1}} \sqrt{c_{n-1}} \psi_{n-1}. \end{cases} \quad (6)$$

with

$$c_n = \frac{e^{2I_0} \theta^2(tV'' + nD + Z)}{\theta(tV'' + (n+1)D + Z) \theta(tV'' + (n-1)D + Z)} \quad (7)$$

$$v_n = -R_0 + \partial_t \ln \frac{\theta(tV'' + (n-1)D + Z)}{\theta(tV'' + nD + Z)}, \quad (8)$$

where  $\theta$  is the classical Riemann theta function.

We know that the degeneracy of solutions of PDE's given in terms of Riemann theta functions gives some important particular solutions. In the case of the NLS equation, we have managed to construct quasi rational solution involving determinant of order  $N$  for each positive integer  $N$  depending on  $2N - 2$  real parameters [13]. In the case of the KdV equation we constructed from Baker Akhiezer functions, solutions in terms of Fredholm determinants and wronskians [54].

In this study, we want to recover Sato formulas for the Toda equation using this method.

From finite gap solutions given in terms of Riemann theta functions, we want to construct by degeneracy, as given in the frame of the NLS equation [79], some quasi-rational solutions and recover the Sato formulas.

Precisely, we derive multisoliton solutions from finite gap solutions by a limit transition, i.e. by making gaps tend toward points in a certain Riemann surface. This will be done in the spirit of the article [79] or more recently [54]. One strength of this approach is that it does not rely on inverse scattering theory or geometric and representation theoretic methods, which offers a fresh perspective on the problem.

We consider the Riemann surface  $\Gamma$  represented by  $\cup_{k=1}^g a_k b_k a_k^{-1} b_k^{-1}$  of the algebraic curve defined by [1, 5]

$$\omega^2 = \prod_{j=1}^{2g+2} (z - E_j),$$

with  $E_j \neq E_k, j \neq k$ .

Let us consider  $\Omega_1''$  and  $\Omega_0'$  [90] abelian integrals verifying

$$\int_{a_k} d\Omega_1''(P) = \int_{a_k} d\Omega_0'(P) = 0 \quad k = 1, \dots, g \quad (9)$$

with the following asymptotic behavior for  $P = (\omega, z) \in \Gamma$  [90],

$$\Omega_1''(P) = \pm \frac{1}{2}(z + R_0 + O(z^{-1})), \quad P \rightarrow P_\infty^\pm \quad (10)$$

$$\Omega_0'(P) = \pm(\ln z - I_0 + O(z^{-1})), \quad P \rightarrow P_\infty^\pm, \quad (11)$$

$$\omega = \pm(z^{g+1} + O(z^g)), \quad P \rightarrow P_\infty^\pm \quad (12)$$

The vectors  $V''$ ,  $D$ ,  $X$  and  $Z$  are defined by

$$V_j'' = \frac{1}{2\pi i} \int_{b_j} d\Omega_1'' \quad (13)$$

$$D = U(P_\infty^+) - U(P_\infty^-), \quad (14)$$

$$X = K + \sum_{j=1}^g U(P_j), \quad Z = U(P_\infty^+) - X, \quad (15)$$

where  $K$  is the vector of Riemann constants.

$U(P)$  is the classical abelian integral  $\int_{P_0}^P dU$ .

The solutions of the system (6) can be written [81, 90],

$$\psi_n(P, t) = \tilde{r}_n(t) e^{n\Omega'_0(P) + t\Omega''_1(P)} \left( \frac{\theta(U(P) + tV'' + nD - X)}{\theta(U(P) - X)} \right), \quad (16)$$

with

$$\tilde{r}_n(t) = r_n(t) \left( \frac{\theta(Z)\theta(Z - D)}{\theta(tV'' + nD + Z)\theta(tV'' + (n-1)D + Z)} \right)^{\frac{1}{2}}, \quad (17)$$

## 2 Degeneracy of solutions

Let us suppose that  $E_j$  are real,  $E_m < E_j$  if  $m < j$  and try to evaluate the limits of all objects in formula (16) when  $E_{2m}, E_{2m-1}$  tends to  $-\alpha_m$ ,  $\alpha_m = \kappa_m^2$ ,  $\kappa_m > 0$ , for  $1 \leq m \leq g$ .

Like in the previous section,  $\theta$  is constructed from the matrix of the B-periods of the surface  $\Gamma$ , the coefficients  $c_{jk}$  being relating with abelian differentials  $dU_j$  by

$$dU_j = \frac{\sum_{k=1}^g c_{jk} z^{g-k}}{\sqrt{\prod_{k=1}^{2g+2} (z - E_k)}} dz, \quad (18)$$

and coefficients  $c_{jk}$  can be obtained by solving the system of linear equations

$$\int_{a_k} dU_j = \delta_{jk}, \quad 1 \leq j \leq g, \quad 1 \leq k \leq g.$$

In the following, we will use the notations:

$$\begin{aligned} \kappa_{mj} &= \sqrt{(\alpha_m + E_{2g+1})(\alpha_j + E_{2g+2})}, & \kappa_{jm} &= \sqrt{(\alpha_j + E_{2g+1})(\alpha_m + E_{2g+2})}, \\ C_k(z) &= \sqrt{(z - E_{2g+1})(\alpha_k - E_{2g+2})}, & D_k(z) &= \sqrt{(z - E_{2g+2})(\alpha_k - E_{2g+1})}, \\ L_k &= \sqrt{\alpha_k + E_{2g+1}}, & M_k &= \sqrt{\alpha_k + E_{2g+2}}, \\ F(z) &= \sqrt{z - E_{2g+1}}, & G(z) &= \sqrt{z - E_{2g+2}}. \end{aligned} \quad (19)$$

### 2.1 Limit of $P(z) = \prod_{j=1}^{2g+2} (z - E_j)$

We study in this paragraph, the limit of  $P(z) = \prod_{j=1}^{2g+2} (z - E_j)$ .

The limit of  $P(z) = \prod_{j=1}^{2g+2} (z - E_j)$  is obviously equal to

$\tilde{P}(z) = \prod_{j=1}^g (z + \alpha_j)^2 (z - E_{2g+1})(z - E_{2g+2})$ ,  
or with (19)

$$\tilde{P}(z) = \prod_{j=1}^g ((z + \alpha_j)F(z)G(z))^2. \quad (20)$$

**2.2 Limit of  $dU_m$**   $= \frac{\sum_{k=1}^g c_{mk} z^{g-k}}{\sqrt{\prod_{k=1}^{2g+2} (z - E_k)}} dz$

In this paragraph, we study the limit of  $dU_m = \frac{\sum_{k=1}^g c_{mk} z^{g-k}}{\sqrt{\prod_{k=1}^{2g+2} (z - E_k)}} dz$ .

The limit of  $dU_m$  is equal to  $d\tilde{U}_m = \frac{\varphi_m(z)}{\prod_{j=1}^g (z + \alpha_j) \sqrt{(z - E_{2g+1})(z - E_{2g+2})}} dz$ ,  
where  $\varphi_m(z) = \sum_{k=1}^g \tilde{c}_{mk} z^{g-k}$ . The normalization condition takes the form in the limit

$$\int_{a_k} dU_j \rightarrow \frac{2\pi i \varphi_j(-\alpha_k)}{\prod_{m \neq k} (\alpha_m - \alpha_k) \sqrt{(\alpha_k + E_{2g+1})(\alpha_k + E_{2g+2})}} = \delta_{kj}, \quad (21)$$

which proves that the numbers  $-\alpha_m$ ,  $m \neq k$  are the zeros of the polynomials  $\varphi_k(z)$ , and so  $\varphi_k(z)$  can be written as  $\varphi_k(z) = \tilde{c}_{k1} \prod_{m \neq k} (z + \alpha_m)$ .  
By (21), we get in the limit

$$\tilde{c}_{k1} = \frac{\sqrt{(\alpha_k + E_{2g+1})(\alpha_k + E_{2g+2})}}{2\pi i}.$$

So

$$\varphi_k(z) = \frac{\sqrt{(\alpha_k + E_{2g+1})(\alpha_k + E_{2g+2})}}{2\pi i} \prod_{m \neq k} (z + \alpha_m).$$

Moreover

$$d\tilde{U}_k(z) = \frac{\varphi_k(z) dz}{\sqrt{(z - E_{2g+1})(z - E_{2g+2})} \prod_{m=1}^g (z + \alpha_m)}$$

in other words

$$d\tilde{U}_k(z) = \frac{\sqrt{(\alpha_k + E_{2g+1})(\alpha_k + E_{2g+2})}}{2\pi i \sqrt{(z - E_{2g+1})(z - E_{2g+2})} (z + \alpha_k)} dz,$$

and with (19)

$$d\tilde{U}_k(z) = \frac{L_k M_k}{2\pi i F(z) G(z) (z + \alpha_k)} dz, \quad (22)$$

### 2.3 Limit of $B_{mk}$

The subject of this subsection is the study of the limit of  $B_{mk}$ .

We have

$$I = \int_{\alpha_m}^{E_{2g+2}} dU_k \rightarrow \frac{1}{2} \tilde{B}_{mk}.$$

The integral  $I$  can be easily evaluate along the real axis on the upper sheet of surface  $\Gamma$  and we get

$$\tilde{B}_{mk} = \frac{i}{\pi} \ln \left| \frac{\sqrt{(\alpha_m + E_{2g+1})(\alpha_k + E_{2g+2})} + \sqrt{(\alpha_k + E_{2g+1})(\alpha_m + E_{2g+2})}}{\sqrt{(\alpha_m + E_{2g+1})(\alpha_k + E_{2g+2})} - \sqrt{(\alpha_k + E_{2g+1})(\alpha_m + E_{2g+2})}} \right|.$$

or with the previous notations (19)

$$\tilde{B}_{mk} = \frac{i}{\pi} \ln \left| \frac{\kappa_{mk} + \kappa_{km}}{\kappa_{mk} - \kappa_{km}} \right|. \quad (23)$$

So  $iB_{kk}$  tends to  $-\infty$ .

Moreover, we have :

$$\int_{E_{2g+2}}^P d\tilde{U}_k = \frac{-i}{2\pi} \ln \left| \frac{\sqrt{(z - E_{2g+1})(\alpha_k + E_{2g+2})} + \sqrt{(z - E_{2g+2})(\alpha_k + E_{2g+1})}}{\sqrt{(z - E_{2g+1})(\alpha_k + E_{2g+2})} - \sqrt{(z - E_{2g+2})(\alpha_k + E_{2g+1})}} \right|.$$

or with the previous notations (19)

$$\int_{E_{2g+2}}^P d\tilde{U}_k = \frac{-i}{2\pi} \ln \left| \frac{C_k(z) + D_k(z)}{C_k(z) - D_k(z)} \right|. \quad (24)$$

### 2.4 Limit of $\Omega_1''$

In this subsection, we study the limit of  $\Omega_1''$ .

$\Omega_1''$  is an abelian integral of the second kind satisfying the conditions

$$\int_{a_k} d\Omega_1''(P) = 0, \quad 1 \leq k \leq g.$$

and such that

$$\Omega_1''(P) = \pm \frac{1}{2}(z + R_0 + O(z^{-1})), \quad P \rightarrow P_\infty^\pm$$

The limit of  $d\Omega_1''(P)$  is equal to  $d\tilde{\Omega}_1''(P) = \frac{\phi(z)}{\prod_{j=1}^g (z + \alpha_j) \sqrt{(z - E_{2g+1})(z - E_{2g+2})}} dz$ ,

where  $\phi(z) = \sum_{k=0}^{g+1} \tilde{c}_k z^{g+1-k}$ .

$\Omega_1''(P)$  satisfying the condition  $\Omega_1''(P) = \pm \frac{1}{2}z + O(1)$  when  $z \rightarrow \pm\infty$ , we have  $\tilde{c}_0 = \frac{1}{2}$ .

Moreover, the conditions

$$\int_{a_k} d\Omega_1''(P) = 0, \quad 1 \leq k \leq g,$$

proves that  $-\alpha_1, \dots, -\alpha_g$  are the zeros of  $\phi$ ;  
thus  $\phi(z) = \frac{1}{2} \prod_{m=1}^g (z + \alpha_m)(z - \beta)$ .

We have,  $P(z) = \sqrt{\prod_{k=1}^{2g+2} (z - E_k)}$  tends to  
 $\tilde{P}(z) = \sqrt{(z - E_{2g+1})(z - E_{2g+2})} \prod_{m=1}^g (z + \alpha_m)$ , we get

$$d\tilde{\Omega}_1''(z) = \frac{z}{2\sqrt{(z - E_{2g+1})(z - E_{2g+2})}} dz,$$

and  $\Omega_1''(P)$  tends to

$$\tilde{\Omega}_1''(P) = \int_{E_{2g+2}}^z \frac{u}{2\sqrt{(u - E_{2g+1})(u - E_{2g+2})}} du.$$

It can be evaluated and it gives

$$\tilde{\Omega}_1''(P) = \frac{E_{2g+1} + E_{2g+2}}{4} \ln \left| \frac{\sqrt{z - E_{2g+1}} + \sqrt{z - E_{2g+2}}}{\sqrt{z - E_{2g+1}} - \sqrt{z - E_{2g+2}}} \right| + \frac{1}{2} \sqrt{(z - E_{2g+1})(z - E_{2g+2})}.$$

With the notations defined in (19), it can be rewritten as

$$\tilde{\Omega}_1''(P) = \frac{E_{2g+1} + E_{2g+2}}{4} \ln \left| \frac{F(z) + G(z)}{F(z) - G(z)} \right| + \frac{1}{2} F(z)G(z). \quad (25)$$

## 2.5 Limit of $\Omega'_0$

We consider  $\Omega'_0$  and we study its limit.

$\Omega'_0$  is an abelian integral of the third kind satisfying the conditions

$$\int_{a_k} d\Omega'_0(P) = 0, \quad 1 \leq k \leq g.$$

and such that

$$\Omega'_0(P) = \pm(\ln z - I_0 + O(z^{-1})), \quad P \rightarrow P_\infty^\pm,$$

The limit of  $d\Omega'_0(P)$  is equal to

$$d\tilde{\Omega}'_0(P) = \frac{\sum_{k=0}^g \tilde{c}_k z^{g-k}}{\prod_{j=1}^g (z + \alpha_j) \sqrt{(z - E_{2g+1})(z - E_{2g+2})}} dz.$$

$\Omega'_0(P)$  satisfying the condition  $\Omega'_0(P) = \pm \ln(z) + O(1)$  when  $z \rightarrow \pm\infty$ , we have  $\tilde{c}_0 = 1$ .

Moreover, the conditions

$$\int_{a_k} d\Omega'_0(P) = 0, \quad 1 \leq k \leq g,$$

prove that  $-\alpha_1, \dots, -\alpha_g$  are the zeros of  $\phi$ ;

thus  $\phi(z)$  defined by  $\phi(z) = \sum_{k=0}^g \tilde{c}_k z^{g-k}$  can be written as



$$\phi(z) = \prod_{m=1}^g (z + \alpha_m).$$

As  $P(z) = \sqrt{\prod_{k=1}^{2g+2} (z - E_k)}$  tends to

$\tilde{P}(z) = \sqrt{(z - E_{2g+1})(z - E_{2g+2})} \prod_{m=1}^g (z + \alpha_m)$ , we get

$$d\tilde{\Omega}'_0(z) = \frac{1}{\sqrt{(z - E_{2g+1})(z - E_{2g+2})}} dz,$$

and  $\Omega'_0(P)$  tends to  $\tilde{\Omega}'_1(P) = \int_{E_{2g+2}}^z \frac{1}{\sqrt{(u - E_{2g+1})(u - E_{2g+2})}} du$ .

It can be calculate and it gives

$$\tilde{\Omega}'_0(P) = \ln \left| \frac{\sqrt{z - E_{2g+1}} + \sqrt{z - E_{2g+2}}}{\sqrt{z - E_{2g+1}} - \sqrt{z - E_{2g+2}}} \right|.$$

It can be written with (19)

$$\tilde{\Omega}'_0(P) = \ln \left| \frac{F(z) + G(z)}{F(z) - G(z)} \right|. \quad (26)$$

## 2.6 Limit of $X$

We deduce the limit of  $X$ .

From the previous sections

$$\tilde{B}_{mk} = \frac{i}{\pi} \ln \left| \frac{\kappa_{mk} + \kappa_{km}}{\kappa_{mk} - \kappa_{km}} \right|,$$

$$\tilde{U}_k(P) = \frac{i}{2\pi} \ln \left| \frac{C_k(z) + D_k(z)}{C_k(z) - D_k(z)} \right|.$$

As  $K$  is defined by  $K_m = \sum_{j=1}^g B_{jm} - \frac{m}{2}$ , and  $X = K + \sum_{j=1}^g U(P_j)$ , we can write

$$X_m \rightarrow \tilde{X}_m = \frac{i}{\pi} \sum_{j=1, j \neq m}^g \ln \left| \frac{\kappa_{mj} + \kappa_{jm}}{\kappa_{mj} - \kappa_{jm}} \right| - \frac{m}{2} + \frac{i}{2\pi} \sum_{j=1}^g \ln \left| \frac{C_m(z_j) + D_m(z_j)}{C_m(z_j) - D_m(z_j)} \right| + i\infty. \quad (27)$$

## 2.7 Limit of $U(P_\infty^+)$ , $U(P_\infty^-)$ , $D$ and $Z$

In this subsection, we study the limit of  $U(P_\infty^+)$ ,  $U(P_\infty^-)$ ,  $D$  and  $Z$ .

From the previous sections, it is easy to get the limits of  $\tilde{U}_k$

$$\lim_{P \rightarrow P_\infty^+} \tilde{U}_k(P) = \frac{-i}{2\pi} \ln \left| \frac{\sqrt{\alpha_k + E_{2g+2}} + \sqrt{\alpha_k + E_{2g+1}}}{\sqrt{\alpha_k + E_{2g+2}} - \sqrt{\alpha_k + E_{2g+1}}} \right|,$$

or

$$\lim_{P \rightarrow P_\infty^+} \tilde{U}_k(P) = \frac{-i}{2\pi} \ln \left| \frac{M_k + L_k}{M_k - L_k} \right|. \quad (28)$$

$$D_k = U_k(P_\infty^+) - U_k(P_\infty^-) \rightarrow \tilde{D}_k = -\frac{i}{2\pi} \ln \frac{M_k + iL_k}{M_k - iL_k}. \quad (29)$$

Thus

$$\lim_{P \rightarrow P_\infty^-} \tilde{U}_k(P) = \frac{-i}{2\pi} \ln \left| \frac{\sqrt{\alpha_k + E_{2g+2}} + \sqrt{\alpha_k + E_{2g+1}}}{\sqrt{\alpha_k + E_{2g+2}} - \sqrt{\alpha_k + E_{2g+1}}} \right| - D_k$$

or

$$\lim_{P \rightarrow P_\infty^-} \tilde{U}_k(P) = \frac{-i}{2\pi} \ln \left| \frac{M_k + L_k}{M_k + L_k} \right| + \frac{i}{2\pi} \ln \left| \frac{M_k + iL_k}{M_k + iL_k} \right|. \quad (30)$$

Therefore the limit of  $Z = U(P_\infty^+) - X$  is given by

$$Z_k \rightarrow \tilde{Z}_k = -\frac{i}{\pi} \sum_{j=1, j \neq k}^g \ln \left| \frac{\kappa_{kj} + \kappa_{jk}}{\kappa_{kj} - \kappa_{jk}} \right| + \frac{k}{2} - \frac{i}{2\pi} \sum_{j=1}^g \ln \left| \frac{C_k(z_j) + D_k(z_j)}{C_k(z_j) - D_k(z_j)} \right| - i\infty - \frac{i}{2\pi} \ln \left| \frac{M_k + L_k}{M_k - L_k} \right| \quad (31)$$

## 2.8 Limit of $V''$

In this paragraph, the limit of  $V''$  is studied.

$V_k'' = \frac{1}{2\pi i} \int_{b_k} d\Omega_1''$  is defined in (13).

We have  $I = \int_{-\alpha_k}^{E_{2g+2}} d\tilde{\Omega}_1'' \rightarrow \frac{1}{2} \int_{b_k} d\tilde{\Omega}_1''$ .

As in the determination of the limit of  $\Omega_1''$ , we have

$$\int_{-\alpha_k}^{E_{2g+2}} d\tilde{\Omega}_1''(P) = -\frac{E_{2g+1} + E_{2g+2}}{4} \ln \left| \frac{\sqrt{\alpha_k + E_{2g+1}} + \sqrt{\alpha_k + E_{2g+2}}}{\sqrt{\alpha_k + E_{2g+1}} - \sqrt{\alpha_k + E_{2g+2}}} \right| - \frac{1}{2} \sqrt{(\alpha_k + E_{2g+1})(\alpha_k + E_{2g+2})},$$

So

$$\begin{aligned} V_k'' &= \frac{1}{2\pi i} \int_{b_k} d\Omega_1'' \rightarrow \tilde{V}'' = -\frac{E_{2g+1} + E_{2g+2}}{4\pi i} \ln \left| \frac{\sqrt{\alpha_k + E_{2g+1}} + \sqrt{\alpha_k + E_{2g+2}}}{\sqrt{\alpha_k + E_{2g+1}} - \sqrt{\alpha_k + E_{2g+2}}} \right| \\ &\quad - \frac{1}{2\pi i} \sqrt{(\alpha_k + E_{2g+1})(\alpha_k + E_{2g+2})}, \\ \tilde{V}_k'' &= \frac{E_{2g+1} + E_{2g+2}}{4\pi i} \ln \left| \frac{L_k - M_k}{L_k + M_k} \right| - \frac{1}{2\pi i} L_k M_k. \end{aligned} \quad (32)$$

## 2.9 Limit of $\theta(p)$

We determine the limit of  $\theta(p)$ .

Let us denote  $A$  the argument of  $\theta(p) = \sum_{k \in \mathbf{Z}^g} \exp\{\pi i(Bk|k) + 2\pi i(k|p)\}$ .

$A$  can be rewritten in the form

$$A = \pi i \sum_{j=1}^g B_{jj} k_j (k_j - 1) + 2\pi i \sum_{j>m} B_{mj} k_m k_j + \sum_{j=1}^g \pi i (2p_j + B_{jj}) k_j.$$

Using the inequality  $k_j(k_j - 1) \geq 0$  for all  $k \in \mathbf{Z}^g$  and the fact that  $iB_{kk}$  tends to  $-\infty$ , we can reduce the limit  $\tilde{\theta}$  of  $\theta(p)$  to a finite sum taken over vectors  $k \in \mathbf{Z}^g$  such that each  $k_j$  must be equal to 0 or 1.

In this section, we compute the limit of all the terms entering in the expression of the solution  $\psi(P, t)$  in  $\theta$ . We will denote  $p_i$  the arguments of these different expressions.

We first study the terms of  $\tilde{r}_n$  corresponding to the arguments  $p_1 = Z = U(P_\infty^+) - X$ ,  $p_2 = Z - D$ ,  $p_3 = tV'' + nD - Z$ ,  $p_4 = tV'' + (n-1)D - Z$ .

Then we will study the remaining terms of  $\psi_n$  corresponding to the arguments  $p_5 = U(P) - X$ ,  $p_6 = U(P) + tV'' + nD - X$ .

With these notations, the solution  $\psi(n, t, z)$  can be written as

$$\psi_n(P, z) = r_n(t) \left( \frac{\theta(p_1)\theta(p_2)}{\theta(p_3)\theta(p_4)} \right)^{\frac{1}{2}} \frac{\theta(p_6)}{\theta(p_5)} e^{n\Omega'_0(P) + t\Omega'_1(P)}.$$

## 2.10 Limit of $\theta(p_1)$

We denote  $p_1$  the term  $Z = U(P_\infty^+) - X$ .

We study its limit.

Then  $\theta(p_1) \rightarrow \tilde{\theta}_1$  :

$$\begin{aligned} \tilde{\theta}_1 = & \sum_{k \in \mathbf{Z}^g, k_j = 0 \text{ or } 1} \exp \left\{ \sum_{j>l} 2 \ln \left| \frac{\kappa_{lj} - \kappa_{jl}}{\kappa_{lj} + \kappa_{jl}} \right| k_j k_l + \sum_{j=1}^g k_j \left( \ln \left| \frac{M_j + L_j}{M_j - L_j} \right| \right. \right. \\ & \left. \left. + \sum_{l \neq j} 2 \ln \left| \frac{\kappa_{jl} + \kappa_{lj}}{\kappa_{jl} - \kappa_{lj}} \right| + j\pi i + \sum_{l=1}^g \ln \left| \frac{C_j(z_l) + D_j(z_l)}{C_j(z_l) - D_j(z_l)} \right| \right) \right\} \end{aligned}$$

## 2.11 Limit of $\theta(p_2)$

We consider  $p_2 = Z - D$ . We study its limit.

Then  $\theta(p_2) \rightarrow \tilde{\theta}_2$  : As  $D_j \rightarrow \tilde{D}_j$ , it is easy to see that

$$\begin{aligned} \tilde{\theta}_2 = & \sum_{k \in \mathbf{Z}^g, k_j = 0 \text{ or } 1} \exp \left\{ \sum_{j>l} 2 \ln \left| \frac{\kappa_{lj} - \kappa_{jl}}{\kappa_{lj} + \kappa_{jl}} \right| k_j k_l + \sum_{j=1}^g k_j \left( \ln \left| \frac{M_j + L_j}{M_j - L_j} \right| \right. \right. \\ & \left. \left. + \sum_{l \neq j} 2 \ln \left| \frac{\kappa_{jl} + \kappa_{lj}}{\kappa_{jl} - \kappa_{lj}} \right| + j\pi i + \sum_{l=1}^g \ln \left| \frac{C_j(z_l) + D_j(z_l)}{C_j(z_l) - D_j(z_l)} \right| + \sum_{j=1}^g \ln \left| \frac{M_j + iL_j}{M_j - iL_j} \right| \right) \right\} \end{aligned}$$

### 2.12 Limit of $\theta(p_3)$

Let  $p_3$  be  $tV'' + nD + Z$ . We study its limit.  
Then  $\theta(p_3) \rightarrow \tilde{\theta}_3$  :

$$\begin{aligned} \tilde{\theta}_3 = & \sum_{k \in \mathbf{Z}^g, k_j = 0 \text{ or } 1} \exp \left\{ \sum_{j>l} 2 \ln \left| \frac{\kappa_{lj} - \kappa_{jl}}{\kappa_{lj} + \kappa_{jl}} \right| k_j k_l + \sum_{j=1}^g k_j \left( \ln \left| \frac{L_j + M_j}{L_j - M_j} \right| \right. \right. \\ & + t \left( \left( \frac{E_{g+1} + E_{2g+2}}{2} \right) \ln \left| \frac{L_j - M_j}{L_j + M_j} \right| - L_j M_j \right) + n \ln \left| \frac{M_j + iL_j}{M_j - iL_j} \right| \\ & \left. \left. + \sum_{l \neq j} 2 \ln \left| \frac{\kappa_{jl} + \kappa_{lj}}{\kappa_{jl} - \kappa_{lj}} \right| + j\pi i + \sum_{l=1}^g \ln \left| \frac{C_j(z_l) + D_j(z_l)}{C_j(z_l) - D_j(z_l)} \right| \right) \right\} \end{aligned}$$

### 2.13 Limit of $\theta(p_4)$

We consider  $p_4 = tV'' + (n-1)D + Z$ ;  $\theta(p_4) \rightarrow \tilde{\theta}_4$ . We determine its limit.  
As  $D_j \rightarrow \tilde{D}_j$ , it is easy to verify that

$$\begin{aligned} \tilde{\theta}_4 = & \sum_{k \in \mathbf{Z}^g, k_j = 0 \text{ or } 1} \exp \left\{ \sum_{j>l} 2 \ln \left| \frac{\kappa_{lj} - \kappa_{jl}}{\kappa_{lj} + \kappa_{jl}} \right| k_j k_l + \sum_{j=1}^g k_j \left( \ln \left| \frac{L_j + M_j}{L_j - M_j} \right| \right. \right. \\ & + t \left( \left( \frac{E_{g+1} + E_{2g+2}}{2} \right) \ln \left| \frac{L_j - M_j}{L_j + M_j} \right| + L_j M_j \right) + (n-1) \ln \left| \frac{M_j + iL_j}{M_j - iL_j} \right| \\ & \left. \left. + \sum_{l \neq j} 2 \ln \left| \frac{\kappa_{jl} + \kappa_{lj}}{\kappa_{jl} - \kappa_{lj}} \right| + j\pi i + \sum_{l=1}^g \ln \left| \frac{C_j(z_l) + D_j(z_l)}{C_j(z_l) - D_j(z_l)} \right| \right) \right\} \end{aligned}$$

### 2.14 Limit of $\theta(p_5)$

The term  $p_5 = U(P) - X$  tends to  $\tilde{p}_5 = \tilde{U}(P) - \tilde{X}$ .

We determine its limit.

Then  $\theta(p_5) \rightarrow \tilde{\theta}_5$  :

$$\begin{aligned} \tilde{\theta}_5 = & \sum_{k \in \mathbf{Z}^g, k_j = 0 \text{ or } 1} \exp \left\{ \sum_{j>l} 2 \ln \left| \frac{\kappa_{lj} - \kappa_{jl}}{\kappa_{lj} + \kappa_{jl}} \right| k_j k_l + \sum_{j=1}^g k_j \left( \ln \left| \frac{C_j(z) - D_j(z)}{C_j(z) + D_j(z)} \right| \right. \right. \\ & \left. \left. + \sum_{l \neq j} 2 \ln \left| \frac{\kappa_{jl} + \kappa_{lj}}{\kappa_{jl} - \kappa_{lj}} \right| + j\pi i + \sum_{l=1}^g \ln \left| \frac{C_j(z_l) + D_j(z_l)}{C_j(z_l) - D_j(z_l)} \right| \right) \right\} \end{aligned}$$

## 2.15 Limit of $\theta(p_6)$

The term  $p_6 = U(P) + tV'' + nD - X$  tends to  $\tilde{p}_6 = \tilde{U}(P) + t\tilde{V}'' - \tilde{X} + n\tilde{D}$ . Then we determine its limit.  $\theta(p_6) \rightarrow \tilde{\theta}_6$  :

$$\begin{aligned} \tilde{\theta}_6 = & \sum_{k \in \mathbf{Z}^g, k_j = 0 \text{ or } 1} \exp \left\{ \sum_{j>l} 2 \ln \left| \frac{\kappa_{lj} - \kappa_{jl}}{\kappa_{lj} + \kappa_{jl}} \right| k_j k_l + \sum_{j=1}^g k_j \left( \ln \left| \frac{L_j + M_j}{L_j - M_j} \right| \right. \right. \\ & + t \left( \left( \frac{E_{g+1} + E_{2g+2}}{2} \right) \ln \left| \frac{L_j - M_j}{L_j + M_j} \right| - L_j M_j \right) + n \ln \left| \frac{M_j + iL_j}{M_j - iL_j} \right| \\ & \left. \left. + \sum_{l \neq j} 2 \ln \left| \frac{\kappa_{jl} + \kappa_{lj}}{\kappa_{jl} - \kappa_{lj}} \right| + j\pi i - \sum_{l=1}^g \ln \left| \frac{C_j(z_l) + D_j(z_l)}{C_j(z_l) - D_j(z_l)} \right| + \ln \left| \frac{C_j(z) - D_j(z)}{C_j(z) + D_j(z)} \right| \right) \right\} \end{aligned}$$

## 2.16 Limit of solutions $\Psi$

From the previous section, we can give the limit  $\tilde{\psi}$  of  $\psi$ . It takes the following form

$$\tilde{\psi}(z, n, t) = \frac{r_n(t) \left( \tilde{\theta}_1 \tilde{\theta}_2 \right)^{\frac{1}{2}}}{\tilde{\theta}_5} \times \frac{e^{\left\{ n \ln \left| \frac{F(z)+G(z)}{F(z)-G(z)} \right| + t \left( \frac{E_{2g+1} + E_{2g+2}}{4} \ln \left| \frac{F(z)+G(z)}{F(z)-G(z)} \right| + \frac{1}{2} F(z)G(z) \right) \right\}} \tilde{\theta}_6}{\left( \tilde{\theta}_3 \tilde{\theta}_4 \right)^{\frac{1}{2}}}$$

In the previous expression,  $\tilde{\theta}_1, \tilde{\theta}_2, \tilde{\theta}_5$  are independent of  $n, t$ .

Only  $\tilde{\theta}_3, \tilde{\theta}_4, \tilde{\theta}_6$  depend on  $n$  and  $t$ .

We choose the particular case in which  $t = 0$ .

We replace  $n$  by  $x$ .

We denote  $c$  the coefficient defined by

$$c = \frac{r_n(0) \left( \tilde{\theta}_1 \tilde{\theta}_2 \right)^{\frac{1}{2}}}{\tilde{\theta}_5}.$$

We denote  $H$  the function defined by

$$H(z) = \left| \frac{F(z) - G(z)}{F(z) + G(z)} \right|$$

In the different sums involving  $\theta$ , only terms with  $k_j = 1$  remain. So the sums can be reduced only on the subsets  $J$  of  $[1; g]$  of integer  $\mathbf{N}$ . We denote  $a_j$  the term  $a_j = -\frac{1}{2} \left| \frac{M_j + iL_j}{M_j - iL_j} \right|$ . Then the term  $\tilde{\theta}_6$  can be written as

$$\tilde{\theta}_6 = \sum_{J \subset \{1, \dots, g\}} \prod_{j, l \in J, j < l} \ln \left| \frac{\kappa_{lj} - \kappa_{jl}}{\kappa_{lj} + \kappa_{jl}} \right|^2 \prod_{j \in J} \left| \frac{L_j + M_j}{L_j - M_j} \right|$$

$$\prod_{l \in J, l \neq j} \ln \left| \frac{\kappa_{jl} + \kappa_{lj}}{\kappa_{jl} - \kappa_{lj}} \right|^2 \prod_{j \in J} \left| \frac{C_j(z_l) + D_j(z_l)}{C_j(z_l) - D_j(z_l)} \right| \left| \frac{C_j(z) - D_j(z)}{C_j(z) + D_j(z)} \right| e^{\sum_{j \in J} j \pi i} e^{-2x \sum_{j \in J} a_j}$$

We denote  $\tau$  the function defined by

$$\tau(x) = \tilde{\theta}_3(x) = \sum_{J \subset \{1, \dots, g\}} \prod_{j, l \in J, j < l} \ln \left| \frac{\kappa_{lj} - \kappa_{jl}}{\kappa_{lj} + \kappa_{jl}} \right|^2 \prod_{j \in J} \left| \frac{L_j + M_j}{L_j - M_j} \right|$$

$$\prod_{l \in J, l \neq j} \ln \left| \frac{\kappa_{jl} + \kappa_{lj}}{\kappa_{jl} - \kappa_{lj}} \right|^2 \prod_{j \in J} \left| \frac{C_j(z_l) + D_j(z_l)}{C_j(z_l) - D_j(z_l)} \right| e^{\sum_{j \in J} j \pi i} e^{-2x \sum_{j \in J} a_j}$$

Thus, the solution  $\tilde{\Psi}$  can be written

$$\tilde{\Psi}(x, z) = c \frac{\exp(xH(z))\chi(x, z)}{\sqrt{\tau(x)\tau(x-1)}}. \quad (33)$$

Here we get back to the Sato formulation of the solution, as given in [94].

This expression is very similar to the solution expressed on page 5830, the expression of the solution is given by

$$\tilde{\Psi}(x, z) = \frac{\exp(xz)\chi(x, z)}{\sqrt{\tau(x)\tau(x-1)}}. \quad (34)$$

The difference between these two statements come from the fact that these two equations are treated differently:

Van Diejen considers the equation

$$a(x)f(x+1) + b(x)f'(x) + a(x-1)f(x-1) = \lambda f(x);$$

here, we consider (??)

$$\frac{r_n}{r_{n+1}} \sqrt{c_n} \psi_{n+1} + v_n \psi_n + \frac{r_n}{r_{n-1}} \sqrt{c_{n-1}} \psi_{n-1} = \lambda \psi_n.$$

## 2.17 Limit of associated potentials

In this subsection, we determine the limit of associated potentials.

The potentials  $u_n = \sqrt{c_n}$  and  $v_n$  are described in (6) and (7)

$$c_n = \frac{e^{2I_0} \theta^2(tV'' + nD + Z)}{\theta(tV'' + (n+1)D + Z) \theta(tV'' + (n-1)D + Z)}$$

$$v_n = -R_0 + \partial_t \ln \frac{\theta(tV'' + (n-1)D + Z)}{\theta(tV'' + nD + Z)}.$$

So the limit of  $u_n$  is equal to

$$\tilde{u}_n = k_1 \frac{\tau(x)}{\sqrt{\tau(x+1)\tau(x-1)}}.$$

which is the same as in [94], (p. 5830) up to the constant  $k_1$ . Denoting  $\sigma(x) = \partial_t \theta(t\tilde{V}'' + x\tilde{D} + \tilde{Z})_{t=0}$ , the limit of  $v_n$  is equal to

$$\tilde{v}_n = k_2 + \frac{\sigma(x-1)}{\tau(x-1)} - \frac{\sigma(x)}{\tau(x)}.$$

which is similar as the one written in [94], (p. 5830) up to the constant  $k_2$ .

### 3 Conclusion

We have used the degeneracy of solutions of some PDEs given in terms of Riemann theta functions to get some important solutions. In particular, in the case of the NLS equation, we have managed to construct quasi rational solution involving determinant of order  $N$  for each positive integer  $N$  depending on  $2N^2$  real parameters [13]. In the case of the KdV equation, from abekian functions, we constructed solutions in terms of Fredholm determinants and wronskians [54].

In this study, we have managed to recover Sato formulas using this method for the Toda equation. From solutions given in terms of Baker-Akiewicz functions, we succeeded to construct by degeneracy, as given in the frame of the NLS equation [79], some quasi-rational solutions and recover the Sato formulas for the Toda equation.

Precisely, we derived multisoliton solutions from finite gap solutions by a limit transition, i.e. by making gaps tend toward points in a certain Riemann surface. One strength of this approach is that it does not rely on inverse scattering theory or geometric and representation theoretic methods, which offers another perspective on the problem.

The degeneracy of the solutions of the NLS equation has allowed to build new quasi-rational solutions of order  $N$  depending on  $2N - 2$  real parameters and their construction up to order 23, previously unknown. In the case of the KdV equation, the degeneracy of the solutions made it possible to find the solutions given by the Darboux method and thus constituted a bridge between the geometric algebra approach and the Darboux transformations framework. In this article, this is somewhat the same situation as that of the KdV equation, where we linked the algebro-geometric approach to the framework of the inverse scattering method.

#### Data Availability

The data that supports the findings of this study are available within the article.

### References

- [1] E.D. Bellokolos, A.I. Bobenko, V.Z. Enol'skii, A.R. Its, V.B. Matveev, Algebro-geometric approach to nonlinear integrable equations, Series in

Nonlinear Dynamics, Springer-Verlag, Berlin, 1996

- [2] E. Date, S. Tanaka, Analogue of inverse Scattering theory for the discrete Hill's equation and exact solutions for the periodic Toda lattice, *Prog. Of Theor. Phys.*, V. 55, N. 2, 1976
- [3] E. Date, S. Tanaka, Periodic Multi-Soliton Solutions of Korteweg-de Vries Equation and Toda Lattice, *Suppl. Of The Prog. Of Theor. Phys.*, N. 59, 1976
- [4] V.N. Duarte, A class of solutions of the two dimensional Toda lattice equation, *Phys. Lett.* V. 385, 126979-1-3, 2021
- [5] B.A. Dubrovin, V.B. Matveev, S.P. Novikov, Integrable systems: non-linear equations of Korteweg-De Vries type, finite-zone linear operators, and abelian varieties, London Mathematical Society, Lecture Notes Series, 60, 53-140, 1976
- [6] B.A. Dubrovin, Theta functions and nonlinear equations, *Russ. Math. Surv.*, V. 36, N. 11, 1-83, 1981
- [7] H. Flaschka, The Toda lattice. II. Existence of integrals, *Phys. Rev. B.*, V. 9, N. 4, 1924-1925, 1974
- [8] P. Gaillard, V.B. Matveev, Wronskian addition formula and its applications, *Max-Planck-Institut für Mathematik, MPI 02-31*, V. 161, 2002
- [9] P. Gaillard, A new family of deformations of Darboux-Pöschl-Teller potentials, *Lett. Math. Phys.*, V. 68, 77-90, 2004
- [10] P. Gaillard, V.B. Matveev, New formulas for the eigenfunctions of the two-particle Calogero-Moser system, *Lett. Math. Phys.*, V. 89, 1-12, 2009
- [11] P. Gaillard, V.B. Matveev, Wronskian and Casorati determinant representations for Darboux-Pöschl-Teller potentials and their difference extensions, *J. Phys A : Math. Theor.*, V. 42, 404409-1-16, 2009
- [12] P. Dubard, P. Gaillard, C. Klein, V. B. Matveev, On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation, *Eur. Phys. J. Spe. Top.*, V. 185, 247-258, 2010
- [13] P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, *J. Phys. A : Meth. Theor.*, V. 44, 435204-1-15, 2011
- [14] P. Gaillard, Wronskian representation of solutions of the NLS equation and higher Peregrine breathers, *J. Math. Sciences : Adv. Appl.*, V. 13, N. 2, 71-153, 2012
- [15] P. Gaillard, Degenerate determinant representation of solution of the NLS equation, higher Peregrine breathers and multi-rogue waves, *J. Math. Phys.*, V. 54, 013504-1-32, 2013



- [16] P. Gaillard, Wronskian representation of solutions of NLS equation and seventh order rogue waves, *J. Mod. Phys.*, V. 4, N. 4, 246-266, 2013
- [17] P. Gaillard, V.B. Matveev, Wronskian addition formula and Darboux-Pöschl-Teller potentials, *J. Math.*, V. 2013, ID 645752, 1-10, 2013
- [18] P. Gaillard, Two-parameters determinant representation of seventh order rogue waves solutions of the NLS equation, *J. Theor. Appl. Phys.*, V. 7, N. 45, 1-6, 2013
- [19] P. Gaillard, Deformations of third order Peregrine breather solutions of the NLS equation with four parameters, *Phys. Rev. E*, V. 88, 042903-1-9, 2013
- [20] P. Gaillard, Two parameters deformations of ninth Peregrine breather solution of the NLS equation and multi rogue waves, *J. Math.*, V. 2013, 1-111, 2013
- [21] P. Gaillard, Six-parameters deformations of fourth order Peregrine breather solutions of the NLS equation, *J. Math. Phys.*, V. 54, 073519-1-22, 2013
- [22] P. Gaillard, The fifth order Peregrine breather and its eight-parameters deformations solutions of the NLS equation, *Commun. Theor. Phys.*, V. 61, 365-369, 2014
- [23] P. Gaillard, Ten parameters deformations of the sixth order Peregrine breather solutions of the NLS equation, *Phys. Scripta*, V. 89, 015004-1-7, 2014
- [24] P. Gaillard, Higher order Peregrine breathers, their deformations and multi-rogue waves, *J. Of Phys. : Conf. Ser.*, V. 482, 012016-1-7, 2014
- [25] P. Gaillard, M. Gastineau, Eighteen parameter deformations of the Peregrine breather of order ten solutions of the NLS equation, *Int. J. Mod. Phys. C*, V. 26, N. 2, 1550016-1-14, 2014
- [26] P. Gaillard, Two parameters wronskian representation of solutions of nonlinear Schrödinger equation, eight Peregrine breather and multi-rogue waves, *J. Math. Phys.*, V. 5, 093506-1-12, 2014
- [27] P. Gaillard, Hierarchy of solutions to the NLS equation and multi-rogue waves, *J. Phys. : Conf. Ser.*, V. 574, 012031-1-5, 2015
- [28] P. Gaillard, Tenth Peregrine breather solution of the NLS, *Ann. Phys.*, V. 355, 293-298, 2015
- [29] P. Gaillard, M. Gastineau, The Peregrine breather of order nine and its deformations with sixteen parameters solutions of the NLS equation *Phys. Lett. A*, V. 379, 1309-1313, 2015

- [30] P. Gaillard, Other  $2N-2$  parameters solutions to the NLS equation and  $2N+1$  highest amplitude of the modulus of the  $N$ -th order AP breather, *J. Phys. A: Math. Theor.*, V. 48, 145203-1-23, 2015
- [31] P. Gaillard, Multi-parametric deformations of the Peregrine breather of order  $N$  solutions to the NLS equation and multi-rogue waves, *Adv. Res.*, V. 4, N. 5, 346-364, 2015
- [32] P. Gaillard, Higher order Peregrine breathers solutions to the NLS equation, *Jour. Phys. : Conf. Ser.*, V. 633, 012106-1-6, 2016
- [33] P. Gaillard, M. Gastineau Patterns of deformations of Peregrine breather of order 3 and 4, solutions to the NLS equation with multi-parameters, *Journal of Theoretical and Applied Physics*, V. 10,1-7, 2016
- [34] P. Gaillard, M. Gastineau Twenty parameters families of solutions to the NLS equation and the eleventh Peregrine breather, *Commun. Theor. Phys*, V. 65, N. 2, 136-144, 2016
- [35] P. Gaillard, Rational solutions to the KPI equation and multi rogue waves, *Annals Of Physics*, V. 367, 1-5, 2016
- [36] P. Gaillard, M. Gastineau Twenty two parameters deformations of the twelfth Peregrine breather solutions to the NLS equation, *Adv. Res.*, V. 10, 83-89, 2016
- [37] P. Gaillard, Towards a classification of the quasi rational solutions to the NLS equation, *Theor. And Math. Phys.*, V. 189, N. 1, 1440-1449, 2016
- [38] P. Gaillard, Fredholm and Wronskian representations of solutions to the KPI equation and multi-rogue waves, *Jour. of Math. Phys.*, V. 57, 063505-1-13, doi: 10.1063/1.4953383, 2016
- [39] P. Gaillard, M. Gastineau Families of deformations of the thirteenth Peregrine breather solutions to the NLS equation depending on twenty four parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. 21, N. 3, 130-139, 2017
- [40] P. Gaillard, From Fredholm and Wronskian representations to rational solutions to the KPI equation depending on  $2N - 2$  parameters, *Int. Jour. of Appl. Sci. And Math.*, V. 4, N. 3, 60-70, 2017
- [41] P. Gaillard, Families of rational solutions of order 5 to the KPI equation depending on 8 parameters, *New Hor. in Math. Phys.*, V. 1, N. 1, 26-31, 2017
- [42] P. Gaillard, 6-th order rational solutions to the KPI Equation depending on 10 parameters, *Jour. Of Bas. And Appl. Res. Int.*, V. 21, N. 2, 92-98, 2017

- [43] P. Gaillard,  $N$ -Order rational solutions to the Johnson equation depending on  $2N - 2$  parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. 4, N. 9, 19-37, 2017
- [44] P. Gaillard, Families of rational solutions to the KPI equation of order 7 depending on 12 parameters, *Int. Jour. of Adv. Res. in Phys. Sci.*, V. 4, N. 11, 24-30, 2017
- [45] P. Gaillard, Rational solutions to the Johnson equation and rogue waves, *Int. Jour. of Inn. In Sci. and Math.*, V. 6, N. 1, 14-19, 2018
- [46] P. Gaillard, Multiparametric families of solutions of the KPI equation, the structure of their rational representations and multi-rogue waves, *Theo. And Mat. Phys.*, V. 196, N. 2, 1174-1199, 2018
- [47] P. Gaillard, The Johnson equation, Fredholm and wronskian representations of solutions and the case of order three, *Adv. In Math. Phys.*, V. 2018, 1-18, 2018
- [48] P. Gaillard, Families of solutions of order 5 to the Johnson equation depending on 8 parameters, *NHIMP*, V. 2, N. 4, 53-61, 2018
- [49] P. Gaillard, Multiparametric families of solutions to the Johnson equation, *J. Phys. : Conf. Series*, V. 1141, 012102-1-10, 2018
- [50] P. Gaillard, Rational solutions to the Boussinesq equation, *Fund. Jour. Of Math. And Appl.*, V. 2, 109-112, 2019
- [51] P. Gaillard, Differential relations for the solutions to the NLS equation and their different representations, *Comm. In Adv. Math. Sci.*, V. 2, N. 4, 1-4, 2019
- [52] P. Gaillard, Multi-parametric families of solutions of order  $N$  to the Boussinesq and KP equations and the degenerate rational case, *UJMA*, V. 3, N. 2, 44-52, 2020
- [53] P. Gaillard, The mKdV equation and multi-parameters rational solutions, *Wave Motion*, V. 100, 102667-1-9, 2021
- [54] P. Gaillard, Degenerate Riemann theta functions, Fredholm and wronskian representations of the solutions to the KdV equation and the degenerate rational case, *Jour. Of Geom. And Phys.*, V. 161, 104059-1-12, 2021
- [55] P. Gaillard,  $2N$  parameter solutions to the Burgers equation, *Jour. Of Appl. Nonlin. Dyn.*, V. 1, N. 1, 1-6, 2021
- [56] P. Gaillard, Multi-Parametric rational Solutions to the KdV equation, *Asi. Jour. Of Res. and Rev. In Phys.*, V. 3, N. 3, 14-21, 2021

- [57] P. Gaillard, Multiparametric solutions to the Gardner equation and the degenerate rational case, *Jour. Of Appl. Ana. And Appl.*, V. 11, N.4, 2102-2113, 2021
- [58] P. Gaillard, Other families of rational solutions to the KPI equation, *Asi. Res. Jour Of Math.*, V. 17, N. 6, 27-34, 2021
- [59] P. Gaillard, Rational solutions to the mKdV equation associated to particular polynomials, *Wave Motion*, V. 107, 102824-1-11, 2021
- [60] P. Gaillard, Multiparametric rational solutions of order  $N$  to the KPI equation and the explicit case of order 3, *Arch. Of cur. Res. Int.*, V. 21, N . 6, 58-71, 2021
- [61] P. Gaillard, Rational solutions to the KPI equation from particular polynomials, *Wave Motion*, V. 108, 102828-1-9, 2022
- [62] P. Gaillard, Rogue Waves of the Lakshmanan Porsezian Daniel Equation Depending on Multi-parameters, *As. Jour. Of Adv. Res. And Rep.*, V 16, N. 3, 32-40, 2022
- [63] P. Gaillard, Quasi-rational and rational solutions to the defocusing NLS equation, *Jour. Of Math. Sci. And Mod. Res. And Rep.*, V 5, N. 1, 24-34, 2022
- [64] P. Gaillard, Rational solutions to the KdV equation in terms of particular polynomials, *Jour. Of Nonlin. Mod. and Ana.*, V. 4, N. 4, 615-626, 2022
- [65] P. Gaillard, Rational solutions to the Painlevé II equation from particular polynomials, *Int. J. Of Appl. Nonlin. Sci.*, V. 3, N. 3, 189-196, 2022
- [66] P. Gaillard, Multi-lump solutions to the KPI equation with a zero degree of derivation, *Jour. Of Theor. And Appl. Phys.*, V. 16, N. 4, 1-8, 2022
- [67] P. Gaillard, Families of solutions to the KPI equation given by an extended Darboux transformation, *Part. Diff. Equ. And Appl.*, V. 3, N. 80, 1-14, 2022
- [68] P. Gaillard, Rogue waves of the Hirota equation in terms of quasi-rational solutions depending on multi-parameters, *WSEAS Trans. On Math.*, DOI: 10.37394/23206.2023.22.24, V. 22, 190-203, 2023
- [69] P. Gaillard, Different representations of the solutions to the cylindrical nonlinear Schrödinger equation, *Dis. Nonlin. And Compl.*, V. 12, N. 3, 539-553, 2023
- [70] P. Gaillard, Rational solutions to the KdV equation from Riemann theta functions, *Jour. Of Theor. And Appl. Phys.*, V. 17, N. 3, 1-5, 2023
- [71] P. Gaillard, On Hadamard's coefficients for two dimensional Schrödinger operators, *Int. Jour. Of Theor. Phys.*, V. 62, N. 205, 1-16, 2023

- [72] P. Gaillard, Rational solutions to the Burgers' equation from particular polynomials, WSEAS Trans. On Math, DOI: 10.37394/23206.2023.22.94, 867-873, 2023
- [73] P. Gaillard, Rational solutions to the Gardner equation from particular polynomials, WSEAS Equ., DOI: 10.37394/232021.2023.3.2, V. 3, 1-24, 2023
- [74] P. Gaillard, Rational solutions to the KPI equation as multi-lumps with a one degree of summation, Int. J. Appl. Comput. Math, doi.org/10.1007/s40819-024-01694-9, 1-14, 2024
- [75] P. Gaillard, Rational solutions to the sixth equation of the NLS hierarchy, ISRG Journal Of Multi. Stud., V. 2, N. 6, 1-14, 2024
- [76] P. Gaillard, Wronskian representation of the solutions to the KdV Equation, Jour. Of App. Phys. Sci. Int., V. 16, N. 2, 1-11, 2024
- [77] P. Gaillard, Quasi-rational solutions to the seventh equation of the NLS hierarchy, Asi. Jour. Of Adv. Res. And Rep., V. 18, N. 7, 256-265, 2024
- [78] P. Gaillard, Solutions of the functional difference Toda equation from centered Darboux transformation, Aequat. Math., 1-21, <https://doi.org/10.1007/s00010-024-01097-7>, 2024
- [79] A.R. Its, A.V. Rybin, M.A. Salle, Exact integration of NLS equation, Theor. Math. Phys., V. 74, N.1, 29-32, 1988
- [80] K. Kajiwara, M. Mazzocco, Y. Ohta, A remark on the Hankel determinan for solutions of thee Toda equation, J. Phys. A: Math. Theor., V. 40, 12661-1675, 2007
- [81] I. Krichever, The periodic non abelian Toda lattice and two dimensional generalization, Uspekhi Math Nauka, V. 36, N.2, 72-77, 1981
- [82] S.V. Manakov, Complete integrability and stochastization of discrete dynamical systems, Zh. Eksp. Teor. Fiz., V. 67, 543-552, 1974
- [83] W.X. Ma, Y. You, Rational solutions of the Toda lattice equation in Casoratian form, Chaos, Solitons And Fractals, V. 22, 395-406, 2004
- [84] W.X. Ma, K.I. Maruno, Complexiton solutions of the Toda lattice, Physica A, V. 343, 219-237, 2004
- [85] V.B. Matveev, Darboux Transformations, Covariance theorems and integrable systems, In L. D. Faddeev's Seminar on Mathematical Physics: American Mathematical Society, 179-209, 2000
- [86] V.B. Matveev, M.A. Salle, Differential-difference evolution equation II, Letters in Mathematical Physics, 3, 425-429, 1979

- [87] A.V. Mikhailov, Integrability of the two-dimensional generalization of the Toda chain, *JETP, Letter*, V. 30, N. 7, 414-418, 1979.
- [88] T. Nilson, C. Schiebold, Solution formulas for the two dimensional Toda lattice and particle like solitons with unexpected asymptotic behaviour, *Jour. Of Nonliner Math. Phys.*, V. 27, N. 1, 57-94, 2020.
- [89] A.O. Smirnov, Finite gap solutions of abelian Toda Chain of genus 4 and 5 in elliptic functions, *Teor. Mat. Fiz*, V. 78, N. 1, 11-21, 1989
- [90] A.O. Smirnov, Elliptic solitons of the integrable nonlinear equations, Thesis, St.-Petersburg State University, 1-170, 2002
- [91] A.A. Stahlhofen, V.B. Matveev, Positons for Toda lattice and related spectral problems, *J. Phys. A: Math. Gen.*, V. 28, 1957-1965, 1995
- [92] Y.L. Sun, W.X. Ma, J.P. Yu, Lump solutions of the 2D equation, *Math. Meth. Appl. Sci.*, 1-7, 2020
- [93] M. Toda, Vibration of a chain with nonlinear interaction, *J. Phys. Soc. Jap.*, V. 6, 22-431, 1967
- [94] J.F. Van Diejen, A Sato formula for reflectionless finite difference operators, *J. of Math. Physics*, V. 40, N. 11, 5822-5834, 1999
- [95] S. Zhang, Y.Y. Zhou, Multiwave solutions for the Toda lattice equation by generalizing exp function method, *Int. Jour. Of Appl. Math.*, V. 44, N. 4, 1-6, 2014